

Appendix A

Gamma and Beta Functions

A.1 A Useful Formula

The following formula is valid:

$$\int_{\mathbf{R}^n} e^{-|x|^2} dx = (\sqrt{\pi})^n.$$

This is an immediate consequence of the corresponding one-dimensional identity

$$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi},$$

which is usually proved from its two-dimensional version by switching to polar coordinates:

$$I^2 = \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{-x^2} e^{-y^2} dy dx = 2\pi \int_0^{\infty} r e^{-r^2} dr = \pi.$$

A.2 Definitions of $\Gamma(z)$ and $B(z, w)$

For a complex number z with $\operatorname{Re} z > 0$ define

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt.$$

$\Gamma(z)$ is called the gamma function. It follows from its definition that $\Gamma(z)$ is analytic on the right half-plane $\operatorname{Re} z > 0$.

Two fundamental properties of the gamma function are that

$$\Gamma(z+1) = z\Gamma(z) \quad \text{and} \quad \Gamma(n) = (n-1)!,$$

where z is a complex number with positive real part and $n \in \mathbf{Z}^+$. Indeed, integration by parts yields

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt = \left[\frac{t^z e^{-t}}{z} \right]_0^\infty + \frac{1}{z} \int_0^\infty t^z e^{-t} dt = \frac{1}{z} \Gamma(z+1).$$

Since $\Gamma(1) = 1$, the property $\Gamma(n) = (n-1)!$ for $n \in \mathbf{Z}^+$ follows by induction. Another important fact is that

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$$

This follows easily from the identity

$$\Gamma\left(\frac{1}{2}\right) = \int_0^\infty t^{-\frac{1}{2}} e^{-t} dt = 2 \int_0^\infty e^{-u^2} du = \sqrt{\pi}.$$

Next we define the beta function. Fix z and w complex numbers with positive real parts. We define

$$B(z, w) = \int_0^1 t^{z-1} (1-t)^{w-1} dt = \int_0^1 t^{w-1} (1-t)^{z-1} dt.$$

We have the following relationship between the gamma and the beta functions:

$$B(z, w) = \frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)},$$

when z and w have positive real parts.

The proof of this fact is as follows:

$$\begin{aligned} \Gamma(z+w)B(z, w) &= \Gamma(z+w) \int_0^1 t^{w-1} (1-t)^{z-1} dt \\ &= \Gamma(z+w) \int_0^\infty u^{w-1} \left(\frac{1}{1+u}\right)^{z+w} du && t = u/(1+u) \\ &= \int_0^\infty \int_0^\infty u^{w-1} \left(\frac{1}{1+u}\right)^{z+w} v^{z+w-1} e^{-v} dv du \\ &= \int_0^\infty \int_0^\infty u^{w-1} s^{z+w-1} e^{-s(u+1)} ds du && s = v/(1+u) \\ &= \int_0^\infty s^z e^{-s} \int_0^\infty (us)^{w-1} e^{-su} du ds \\ &= \int_0^\infty s^{z-1} e^{-s} \Gamma(w) ds \\ &= \Gamma(z)\Gamma(w). \end{aligned}$$

A.3 Volume of the Unit Ball and Surface of the Unit Sphere

We denote by v_n the volume of the unit ball in \mathbf{R}^n and by ω_{n-1} the surface area of the unit sphere \mathbf{S}^{n-1} . We have the following:

$$\omega_{n-1} = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$$

and

$$v_n = \frac{\omega_{n-1}}{n} = \frac{2\pi^{\frac{n}{2}}}{n\Gamma(\frac{n}{2})} = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2} + 1)}.$$

The easy proofs are based on the formula in Appendix A.1. We have

$$(\sqrt{\pi})^n = \int_{\mathbf{R}^n} e^{-|x|^2} dx = \omega_{n-1} \int_0^\infty e^{-r^2} r^{n-1} dr,$$

by switching to polar coordinates. Now change variables $t = r^2$ to obtain that

$$\pi^{\frac{n}{2}} = \frac{\omega_{n-1}}{2} \int_0^\infty e^{-t} t^{\frac{n}{2}-1} dt = \frac{\omega_{n-1}}{2} \Gamma\left(\frac{n}{2}\right).$$

This proves the formula for the surface area of the unit sphere in \mathbf{R}^n .

To compute v_n , write again using polar coordinates

$$v_n = |B(0, 1)| = \int_{|x| \leq 1} 1 dx = \int_{\mathbf{S}^{n-1}} \int_0^1 r^{n-1} dr d\theta = \frac{1}{n} \omega_{n-1}.$$

Here is another way to relate the volume to the surface area. Let $B(0, R)$ be the ball in \mathbf{R}^n of radius $R > 0$ centered at the origin. Then the volume of the shell $B(0, R+h) \setminus B(0, R)$ divided by h tends to the surface area of $B(0, R)$ as $h \rightarrow 0$. In other words, the derivative of the volume of $B(0, R)$ with respect to the radius R is equal to the surface area of $B(0, R)$. Since the volume of $B(0, R)$ is $v_n R^n$, it follows that the surface area of $B(0, R)$ is $n v_n R^{n-1}$. Taking $R = 1$, we deduce $\omega_{n-1} = n v_n$.

A.4 Computation of Integrals Using Gamma Functions

Let k_1, \dots, k_n be nonnegative even integers. The integral

$$\int_{\mathbf{R}^n} x_1^{k_1} \dots x_n^{k_n} e^{-|x|^2} dx_1 \dots dx_n = \prod_{j=1}^n \int_{-\infty}^{+\infty} x_j^{k_j} e^{-x_j^2} dx_j = \prod_{j=1}^n \Gamma\left(\frac{k_j+1}{2}\right)$$

expressed in polar coordinates is equal to

$$\left(\int_{\mathbf{S}^{n-1}} \theta_1^{k_1} \dots \theta_n^{k_n} d\theta \right) \int_0^\infty r^{k_1+\dots+k_n} r^{n-1} e^{-r^2} dr,$$

where $\theta = (\theta_1, \dots, \theta_n)$. This leads to the identity

$$\int_{\mathbb{S}^{n-1}} \theta_1^{k_1} \dots \theta_n^{k_n} d\theta = 2\Gamma\left(\frac{k_1 + \dots + k_n + n}{2}\right)^{-1} \prod_{j=1}^n \Gamma\left(\frac{k_j + 1}{2}\right).$$

Another classical integral that can be computed using gamma functions is the following:

$$\int_0^{\pi/2} (\sin \varphi)^a (\cos \varphi)^b d\varphi = \frac{1}{2} \frac{\Gamma(\frac{a+1}{2})\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+b+2}{2})},$$

whenever a and b are complex numbers with $\operatorname{Re} a > -1$ and $\operatorname{Re} b > -1$.

Indeed, change variables $u = (\sin \varphi)^2$; then $du = 2(\sin \varphi)(\cos \varphi)d\varphi$, and the preceding integral becomes

$$\frac{1}{2} \int_0^1 u^{\frac{a-1}{2}} (1-u)^{\frac{b-1}{2}} du = \frac{1}{2} B\left(\frac{a+1}{2}, \frac{b+1}{2}\right) = \frac{1}{2} \frac{\Gamma(\frac{a+1}{2})\Gamma(\frac{b+1}{2})}{\Gamma(\frac{a+b+2}{2})}.$$

A.5 Meromorphic Extensions of $B(z, w)$ and $\Gamma(z)$

Using the identity $\Gamma(z+1) = z\Gamma(z)$, we can easily define a meromorphic extension of the gamma function on the whole complex plane starting from its known values on the right half-plane. We give an explicit description of the meromorphic extension of $\Gamma(z)$ on the whole plane. First write

$$\Gamma(z) = \int_0^1 t^{z-1} e^{-t} dt + \int_1^\infty t^{z-1} e^{-t} dt$$

and observe that the second integral is an analytic function of z for all $z \in \mathbf{C}$. Write the first integral as

$$\int_0^1 t^{z-1} \left\{ e^{-t} - \sum_{j=0}^N \frac{(-t)^j}{j!} \right\} dt + \sum_{j=0}^N \frac{(-1)^j / j!}{z+j}.$$

The last integral converges when $\operatorname{Re} z > -N-1$, since the expression inside the curly brackets is $O(t^{N+1})$ as $t \rightarrow 0$. It follows that the gamma function can be defined to be an analytic function on $\operatorname{Re} z > -N-1$ except at the points $z = -j$, $j = 0, 1, \dots, N$, at which it has simple poles with residues $\frac{(-1)^j}{j!}$. Since N was arbitrary, it follows that the gamma function has a meromorphic extension on the whole plane.

In view of the identity

$$B(z, w) = \frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)},$$

the definition of $B(z, w)$ can be extended to $\mathbf{C} \times \mathbf{C}$. It follows that $B(z, w)$ is a meromorphic function in each argument.

A.6 Asymptotics of $\Gamma(x)$ as $x \rightarrow \infty$

We now derive *Stirling's formula*:

$$\lim_{x \rightarrow \infty} \frac{\Gamma(x+1)}{\left(\frac{x}{e}\right)^x \sqrt{2\pi x}} = 1.$$

First change variables $t = x + sx\sqrt{\frac{2}{x}}$ to obtain

$$\Gamma(x+1) = \int_0^\infty e^{-t} t^x dt = \left(\frac{x}{e}\right)^x \sqrt{2x} \int_{-\sqrt{x/2}}^{+\infty} \frac{\left(1 + s\sqrt{\frac{2}{x}}\right)^x}{e^{2s\sqrt{x/2}}} ds.$$

Setting $y = \sqrt{\frac{x}{2}}$, we obtain

$$\frac{\Gamma(x+1)}{\left(\frac{x}{e}\right)^x \sqrt{2x}} = \int_{-\infty}^{+\infty} \left(\frac{(1 + \frac{s}{y})^y}{e^s}\right)^{2y} \chi_{(-y, \infty)}(s) ds.$$

To show that the last integral converges to $\sqrt{\pi}$ as $y \rightarrow \infty$, we need the following:

(1) The fact that

$$\lim_{y \rightarrow \infty} \left(\frac{(1 + s/y)^y}{e^s}\right)^{2y} \rightarrow e^{-s^2},$$

which follows easily by taking logarithms and applying L'Hôpital's rule twice.

(2) The estimate, valid for $y \geq 1$,

$$\left(\frac{(1 + \frac{s}{y})^y}{e^s}\right)^{2y} \leq \begin{cases} \frac{(1+s)^2}{e^s} & \text{when } s \geq 0, \\ e^{-s^2} & \text{when } -y < s < 0, \end{cases}$$

which can be easily checked using calculus. Using these facts, the Lebesgue dominated convergence theorem, the trivial fact that $\chi_{-y < s < \infty} \rightarrow 1$ as $y \rightarrow \infty$, and the identity in Appendix A.1, we obtain that

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{\Gamma(x+1)}{\left(\frac{x}{e}\right)^x \sqrt{2x}} &= \lim_{y \rightarrow \infty} \int_{-\infty}^{+\infty} \left(\frac{(1 + \frac{s}{y})^y}{e^s}\right)^{2y} \chi_{(-y, \infty)}(s) ds \\ &= \int_{-\infty}^{+\infty} e^{-s^2} ds \\ &= \sqrt{\pi}. \end{aligned}$$

As a consequence of Stirling's formula, for any $t > 0$, we obtain

$$\lim_{x \rightarrow \infty} \frac{\Gamma(x)}{\Gamma(x+t)} = 0.$$

A.7 Euler's Limit Formula for the Gamma Function

For n a positive integer and $\operatorname{Re} z > 0$ we consider the functions

$$\Gamma_n(z) = \int_0^n \left(1 - \frac{t}{n}\right)^n t^{z-1} dt$$

We show that

$$\Gamma_n(z) = \frac{n! n^z}{z(z+1)\cdots(z+n)}$$

and we obtain *Euler's limit formula for the gamma function*

$$\lim_{n \rightarrow \infty} \Gamma_n(z) = \Gamma(z).$$

We write $\Gamma(z) - \Gamma_n(z) = I_1(z) + I_2(z) + I_3(z)$, where

$$\begin{aligned} I_1(z) &= \int_n^\infty e^{-t} t^{z-1} dt, \\ I_2(z) &= \int_{n/2}^n \left(e^{-t} - \left(1 - \frac{t}{n}\right)^n \right) t^{z-1} dt, \\ I_3(z) &= \int_0^{n/2} \left(e^{-t} - \left(1 - \frac{t}{n}\right)^n \right) t^{z-1} dt. \end{aligned}$$

Obviously $I_1(z)$ tends to zero as $n \rightarrow \infty$. For I_2 and I_3 we have that $0 \leq t < n$, and by the Taylor expansion of the logarithm we obtain

$$\log \left(1 - \frac{t}{n}\right)^n = n \log \left(1 - \frac{t}{n}\right) = -t - L,$$

where

$$L = \frac{t^2}{n} \left(\frac{1}{2} + \frac{1}{3} \frac{t}{n} + \frac{1}{4} \frac{t^2}{n^2} + \cdots \right).$$

It follows that

$$0 < e^{-t} - \left(1 - \frac{t}{n}\right)^n = e^{-t} - e^{-L} e^{-t} \leq e^{-t},$$

and thus $I_2(z)$ tends to zero as $n \rightarrow \infty$. For I_3 we have $t/n \leq 1/2$, which implies that

$$L \leq \frac{t^2}{n} \sum_{k=0}^{\infty} \frac{1}{(k+1)2^{k-1}} = \frac{t^2}{n} c.$$

Consequently, for $t/n \leq 1/2$ we have

$$0 \leq e^{-t} - \left(1 - \frac{t}{n}\right)^n = e^{-t}(1 - e^{-L}) \leq e^{-t}L \leq e^{-t} \frac{ct^2}{n}.$$

Plugging this estimate into I_3 , we deduce that

$$|I_3(z)| \leq \frac{c}{n} \Gamma(\operatorname{Re} z + 2),$$

which certainly tends to zero as $n \rightarrow \infty$.

Next, n integrations by parts give

$$\Gamma_n(z) = \frac{n}{nz} \frac{n-1}{n(z+1)} \frac{n-2}{n(z+2)} \cdots \frac{1}{n(z+n-1)} \int_0^n t^{z+n-1} dt = \frac{n! n^z}{z(z+1) \cdots (z+n)}.$$

This can be written as

$$1 = \Gamma_n(z) z \exp \left\{ z \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n \right) \right\} \prod_{k=1}^n \left(1 + \frac{z}{k} \right) e^{-z/k}.$$

Taking limits as $n \rightarrow \infty$, we obtain an *infinite product form of Euler's limit formula*,

$$1 = \Gamma(z) z e^{\gamma z} \prod_{k=1}^{\infty} \left(1 + \frac{z}{k} \right) e^{-z/k},$$

where $\operatorname{Re} z > 0$ and γ is *Euler's constant*

$$\gamma = \lim_{n \rightarrow \infty} 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n.$$

The infinite product converges uniformly on compact subsets of the complex plane that excludes $z = 0, -1, -2, \dots$, and thus it represents a holomorphic function in this domain. This holomorphic function multiplied by $\Gamma(z) z e^{\gamma z}$ is equal to 1 on $\operatorname{Re} z > 0$ and by analytic continuation it must be equal to 1 on $\mathbf{C} \setminus \{0, -1, -2, \dots\}$. But $\Gamma(z)$ has simple poles, while the infinite product vanishes to order one, at the nonpositive integers. We conclude that Euler's limit formula holds for all complex numbers z ; consequently, $\Gamma(z)$ has no zeros and $\Gamma(z)^{-1}$ is entire.

An immediate consequence of Euler's limit formula is the identity

$$\frac{1}{|\Gamma(x+iy)|^2} = \frac{1}{|\Gamma(x)|^2} \prod_{k=0}^{\infty} \left(1 + \frac{y^2}{(k+x)^2} \right),$$

which holds for x and y real with $x \notin \{0, -1, -2, \dots\}$. As a consequence we have that

$$|\Gamma(x+iy)| \leq |\Gamma(x)|$$

and also that

$$\frac{1}{|\Gamma(x+iy)|} \leq \frac{1}{|\Gamma(x)|} e^{C(x)|y|^2},$$

where

$$C(x) = \frac{1}{2} \sum_{k=0}^{\infty} \frac{1}{(k+x)^2},$$

whenever $y \in \mathbf{R}$ and $x \in \mathbf{R} \setminus \{0, -1, -2, \dots\}$. Before we find a similar estimate for $x \in \{0, -1, -2, \dots\}$ we provide a simpler expression for this estimate when $x > 0$.

When $x > 0$ we have

$$C(x) \leq \frac{1}{2x^2} + \frac{1}{2} \sum_{k=1}^{\infty} \frac{1}{(k+x)^2} \leq \frac{1}{2x^2} + \frac{1}{2} \int_0^{\infty} \frac{dt}{(t+x)^2} = \frac{1}{2x^2} + \frac{1}{2x} \int_1^{\infty} \frac{dt}{t^2} = \frac{1}{2x^2} + \frac{1}{2x}.$$

Thus we conclude that when $x > 0$ and $y \in \mathbf{R}$ we have

$$\frac{1}{|\Gamma(x+iy)|} \leq \frac{1}{|\Gamma(x)|} e^{\max\{x^{-2}, x^{-1}\}|y|^2}.$$

When $x = 0$ we write $\Gamma(iy)iy = \Gamma(1+iy)$ and use the preceding inequality to obtain

$$\frac{1}{|\Gamma(iy)|} \leq \frac{|iy|}{|\Gamma(1)|} e^{|y|^2} = |y|e^{|y|^2}$$

and more generally for $x = -N \in \{-1, -2, \dots\}$ and $y \in \mathbf{R}$ we obtain by induction

$$\frac{1}{|\Gamma(-N+iy)|} \leq |iy| |1+iy| |2+iy| \cdots |N+iy| e^{|y|^2}.$$

A.8 Reflection and Duplication Formulas for the Gamma Function

The *reflection formula* relates the values of the gamma function of a complex number z and its reflection about the point $1/2$ in the following way:

$$\frac{\sin(\pi z)}{\pi} = \frac{1}{\Gamma(z)} \frac{1}{\Gamma(1-z)}.$$

The *duplication formula* relates the entire functions $\Gamma(2z)^{-1}$ and $\Gamma(z)^{-1}$ as follows:

$$\frac{1}{\Gamma(z)\Gamma(z+\frac{1}{2})} = \frac{\pi^{-\frac{1}{2}} 2^{2z-1}}{\Gamma(2z)}.$$

Both of these could be proved using Euler's limit formula. The reflection formula also uses the identity

$$\prod_{k=1}^{\infty} \left(1 - \frac{z^2}{k^2}\right) = \frac{\sin(\pi z)}{\pi z},$$

while the duplication formula makes use of the fact that

$$\lim_{n \rightarrow \infty} \frac{(n!)^2 2^{2n+1}}{(2n)! n^{1/2}} = 2\pi^{1/2}.$$

These and other facts related to the gamma function can be found in Olver [271].

Appendix B

Bessel Functions

B.1 Definition

We survey some basics from the theory of Bessel functions J_ν of complex order ν with $\operatorname{Re} \nu > -1/2$. We define the Bessel function J_ν of order ν by its *Poisson representation formula*

$$J_\nu(t) = \frac{\left(\frac{t}{2}\right)^\nu}{\Gamma(\nu + \frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^{+1} e^{its} (1-s^2)^\nu \frac{ds}{\sqrt{1-s^2}},$$

where $\operatorname{Re} \nu > -1/2$ and $t \geq 0$. Although this definition is also valid when t is a complex number, for the applications we have in mind, it suffices to consider the case that t is real and nonnegative; in this case $J_\nu(t)$ is also a real number.

B.2 Some Basic Properties

Let us summarize a few properties of Bessel functions. We take $t > 0$.

(1) We have the following recurrence formula:

$$\frac{d}{dt} (t^{-\nu} J_\nu(t)) = -t^{-\nu} J_{\nu+1}(t), \quad \operatorname{Re} \nu > -1/2.$$

(2) We also have the companion recurrence formula:

$$\frac{d}{dt} (t^\nu J_\nu(t)) = t^\nu J_{\nu-1}(t), \quad \operatorname{Re} \nu > 1/2.$$

(3) $J_\nu(t)$ satisfies the differential equation:

$$t^2 \frac{d^2}{dt^2}(J_\nu(t)) + t \frac{d}{dt}(J_\nu(t)) + (t^2 - \nu^2)J_\nu(t) = 0.$$

(4) If $\nu \in \mathbf{Z}^+$, then we have the following identity, which was taken by Bessel as the definition of J_ν for integer ν :

$$J_\nu(t) = \frac{1}{2\pi} \int_0^{2\pi} e^{it \sin \theta} e^{-i\nu \theta} d\theta = \frac{1}{2\pi} \int_0^{2\pi} \cos(t \sin \theta - \nu \theta) d\theta.$$

(5) For $\operatorname{Re} \nu > -1/2$ we have the following identity:

$$J_\nu(t) = \frac{1}{\Gamma(\frac{1}{2})} \left(\frac{t}{2}\right)^\nu \sum_{j=0}^{\infty} (-1)^j \frac{\Gamma(j + \frac{1}{2})}{\Gamma(j + \nu + 1)} \frac{t^{2j}}{(2j)!},$$

which can also be written as

$$J_\nu(t) = \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \frac{1}{\Gamma(j + \nu + 1)} \left(\frac{t}{2}\right)^{2j+\nu},$$

using the equality $(2j)! = 2^{2j} j! (j - \frac{1}{2})(j - \frac{3}{2}) \cdots \frac{1}{2} = 2^{2j} j! \Gamma(j + \frac{1}{2}) \Gamma(\frac{1}{2})^{-1}$.

(6) For $\operatorname{Re} \nu > 1/2$ the identity below is valid:

$$\frac{d}{dt}(J_\nu(t)) = \frac{1}{2}(J_{\nu-1}(t) - J_{\nu+1}(t)).$$

We first verify property (1). We have

$$\begin{aligned} \frac{d}{dt}(t^{-\nu} J_\nu(t)) &= \frac{i}{2^\nu \Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 s e^{its} (1-s^2)^{\nu-\frac{1}{2}} ds \\ &= \frac{i}{2^\nu \Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 \frac{it}{2} e^{its} \frac{(1-s^2)^{\nu+\frac{1}{2}}}{\nu + \frac{1}{2}} ds \\ &= -t^{-\nu} J_{\nu+1}(t), \end{aligned}$$

where we integrated by parts and used the fact that $\Gamma(x+1) = x\Gamma(x)$.

We now prove Property (2) for $\operatorname{Re} \nu > 1/2$:

$$\begin{aligned} &\frac{d}{dt}(t^\nu J_\nu(t)) \\ &= \frac{2\nu t^{2\nu-1} 2^{-\nu}}{\Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} (1-s^2)^{\nu-\frac{1}{2}} ds + \frac{t^{2\nu} 2^{-\nu}}{\Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2})} i \int_{-1}^1 e^{its} is (1-s^2)^{\nu-\frac{1}{2}} ds \end{aligned}$$

$$\begin{aligned}
 &= \frac{2vt^{2v-1}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its}(1-s^2)^{v-\frac{1}{2}} ds + \frac{t^{2v}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 \left(\frac{e^{its}}{t}\right)'(1-s^2)^{v-\frac{1}{2}} s ds \\
 &= \frac{2vt^{2v-1}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its}(1-s^2)^{v-\frac{1}{2}} ds - \frac{t^{2v}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 \frac{e^{its}}{t} ((1-s^2)^{v-\frac{1}{2}} s)' ds \\
 &= \frac{t^{2v-1}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} \left[2v(1-s^2)^{v-\frac{1}{2}} - ((1-s^2)^{v-\frac{1}{2}} s)' \right] ds \\
 &= \frac{t^{2v-1}2^{-v}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} (2v-1)(1-s^2)^{v-\frac{3}{2}} ds \\
 &= \frac{t^{2v-1}2^{-(v-1)}}{\Gamma(v-\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} (1-s^2)^{v-\frac{1}{2}} \frac{ds}{\sqrt{1-s^2}} \\
 &= t^v J_{v-1}(t).
 \end{aligned}$$

We proceed with the proof of property (3). A calculation using the definition of the Bessel function gives that the left-hand side of (3) is equal to

$$\frac{2^{-v}t^{v+1}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^{+1} e^{ist} \left((1-s^2)t + 2is(v+\frac{1}{2}) \right) (1-s^2)^{v-\frac{1}{2}} ds,$$

which in turn is equal to

$$-i \frac{2^{-v}t^{v+1}}{\Gamma(v+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^{+1} \frac{d}{ds} \left(e^{ist} (1-s^2)^{v+\frac{1}{2}} \right) ds = 0.$$

Property (4) can be derived directly from (1). Define

$$G_v(t) = \frac{1}{2\pi} \int_0^{2\pi} e^{it \sin \theta} e^{-iv\theta} d\theta,$$

for $v = 0, 1, 2, \dots$ and $t > 0$. We can show easily that $G_0 = J_0$. If we had

$$\frac{d}{dt} (t^{-v} G_v(t)) = -t^{-v} G_{v+1}(t), \quad t > 0,$$

for $v \in \mathbf{Z}^+$, we would immediately conclude that $G_v = J_v$ for $v \in \mathbf{Z}^+$. We have

$$\begin{aligned}
 \frac{d}{dt} (t^{-v} G_v(t)) &= -t^{-v} \left(\frac{v}{t} G_v(t) - \frac{dG_v}{dt}(t) \right) \\
 &= -t^{-v} \int_0^{2\pi} \frac{v}{2\pi t} e^{it \sin \theta} e^{-iv\theta} - \frac{1}{2\pi} \left(\frac{d}{dt} e^{it \sin \theta} \right) e^{-iv\theta} d\theta \\
 &= -\frac{t^{-v}}{2\pi} \int_0^{2\pi} i \frac{d}{d\theta} \left(\frac{e^{it \sin \theta - iv\theta}}{t} \right) + (\cos \theta - i \sin \theta) e^{it \sin \theta} e^{-iv\theta} d\theta \\
 &= -\frac{t^{-v}}{2\pi} \int_0^{2\pi} e^{it \sin \theta} e^{-i(v+1)\theta} d\theta \\
 &= -t^{-v} G_{v+1}(t).
 \end{aligned}$$

For t real, the identity in (5) can be derived by inserting the expression

$$\sum_{j=0}^{\infty} (-1)^j \frac{(ts)^{2j}}{(2j)!} + i \sin(ts)$$

for e^{its} in the definition of the Bessel function $J_\nu(t)$ in Appendix B.1. Algebraic manipulations yield

$$\begin{aligned} J_\nu(t) &= \frac{(t/2)^\nu}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} (-1)^j \frac{1}{\Gamma(\nu + \frac{1}{2})} \frac{t^{2j}}{(2j)!} 2 \int_0^1 s^{2j-1} (1-s^2)^{\nu-\frac{1}{2}} s ds \\ &= \frac{(t/2)^\nu}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} (-1)^j \frac{1}{\Gamma(\nu + \frac{1}{2})} \frac{t^{2j}}{(2j)!} \frac{\Gamma(j + \frac{1}{2})\Gamma(\nu + \frac{1}{2})}{\Gamma(j + \nu + 1)} \\ &= \frac{(t/2)^\nu}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} (-1)^j \frac{\Gamma(j + \frac{1}{2})}{\Gamma(j + \nu + 1)} \frac{t^{2j}}{(2j)!}. \end{aligned}$$

To derive property (6) we first multiply (1) by t^ν and (2) by $t^{-\nu}$; then we use the product rule for differentiation and we add the resulting expressions.

For further identities on Bessel functions, one may consult Watson's monograph [371].

B.3 An Interesting Identity

Let $\operatorname{Re} \mu > -\frac{1}{2}$, $\operatorname{Re} \nu > -1$, and $t > 0$. Then the following identity is valid:

$$\int_0^1 J_\mu(ts) s^{\mu+1} (1-s^2)^\nu ds = \frac{\Gamma(\nu+1)2^\nu}{t^{\nu+1}} J_{\mu+\nu+1}(t).$$

To prove this identity we use formula (5) in Appendix B.2. We have

$$\begin{aligned} &\int_0^1 J_\mu(ts) s^{\mu+1} (1-s^2)^\nu ds \\ &= \frac{(\frac{t}{2})^\mu}{\Gamma(\frac{1}{2})} \int_0^1 \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(j + \frac{1}{2}) t^{2j}}{\Gamma(j + \mu + 1) (2j)!} s^{2j+\mu+\mu} (1-s^2)^\nu s ds \\ &= \frac{1}{2} \frac{(\frac{t}{2})^\mu}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(j + \frac{1}{2}) t^{2j}}{\Gamma(j + \mu + 1) (2j)!} \int_0^1 u^{j+\mu} (1-u)^\nu du \\ &= \frac{1}{2} \frac{(\frac{t}{2})^\mu}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(j + \frac{1}{2}) t^{2j}}{\Gamma(j + \mu + 1) (2j)!} \frac{\Gamma(\mu + j + 1) \Gamma(\nu + 1)}{\Gamma(\mu + \nu + j + 2)} \\ &= \frac{2^\nu \Gamma(\nu + 1)}{t^{\nu+1}} \frac{(\frac{t}{2})^{\mu+\nu+1}}{\Gamma(\frac{1}{2})} \sum_{j=0}^{\infty} \frac{(-1)^j \Gamma(j + \frac{1}{2}) t^{2j}}{\Gamma(j + \mu + \nu + 2) (2j)!} \\ &= \frac{\Gamma(\nu + 1) 2^\nu}{t^{\nu+1}} J_{\mu+\nu+1}(t). \end{aligned}$$

B.4 The Fourier Transform of Surface Measure on \mathbf{S}^{n-1}

Let $d\sigma$ denote surface measure on \mathbf{S}^{n-1} for $n \geq 2$. Then the following is true:

$$\widehat{d\sigma}(\xi) = \int_{\mathbf{S}^{n-1}} e^{-2\pi i \xi \cdot \theta} d\theta = \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} J_{\frac{n-2}{2}}(2\pi|\xi|).$$

To see this, use the result in Appendix D.3 to write

$$\begin{aligned} \widehat{d\sigma}(\xi) &= \int_{\mathbf{S}^{n-1}} e^{-2\pi i \xi \cdot \theta} d\theta \\ &= \frac{2\pi^{\frac{n-1}{2}}}{\Gamma(\frac{n-1}{2})} \int_{-1}^{+1} e^{-2\pi i |\xi| s} (1-s^2)^{\frac{n-2}{2}} \frac{ds}{\sqrt{1-s^2}} \\ &= \frac{2\pi^{\frac{n-1}{2}}}{\Gamma(\frac{n-1}{2})} \frac{\Gamma(\frac{n-2}{2} + \frac{1}{2}) \Gamma(\frac{1}{2})}{(\pi|\xi|)^{\frac{n-2}{2}}} J_{\frac{n-2}{2}}(2\pi|\xi|) \\ &= \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} J_{\frac{n-2}{2}}(2\pi|\xi|). \end{aligned}$$

B.5 The Fourier Transform of a Radial Function on \mathbf{R}^n

Let $f(x) = f_0(|x|)$ be a radial function defined on \mathbf{R}^n , where f_0 is defined on $[0, \infty)$. Then the Fourier transform of f is given by the formula

$$\widehat{f}(\xi) = \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} \int_0^\infty f_0(r) J_{\frac{n}{2}-1}(2\pi r|\xi|) r^{\frac{n}{2}} dr.$$

To obtain this formula, use polar coordinates to write

$$\begin{aligned} \widehat{f}(\xi) &= \int_{\mathbf{R}^n} f(x) e^{-2\pi i \xi \cdot x} dx \\ &= \int_0^\infty \int_{\mathbf{S}^{n-1}} f_0(r) e^{-2\pi i \xi \cdot r\theta} d\theta r^{n-1} dr \\ &= \int_0^\infty f_0(r) \widehat{d\sigma}(r\xi) r^{n-1} dr \\ &= \int_0^\infty f_0(r) \frac{2\pi}{(r|\xi|)^{\frac{n-2}{2}}} J_{\frac{n-2}{2}}(2\pi r|\xi|) r^{n-1} dr \\ &= \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} \int_0^\infty f_0(r) J_{\frac{n}{2}-1}(2\pi r|\xi|) r^{\frac{n}{2}} dr. \end{aligned}$$

As an application we take $f(x) = \chi_{B(0,1)}(x)$, where $B(0,1)$ is the unit ball in \mathbf{R}^n . We obtain

$$(\chi_{B(0,1)})^\wedge(\xi) = \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} \int_0^1 J_{\frac{n}{2}-1}(2\pi|\xi|r)r^{\frac{n}{2}} dr = \frac{J_{\frac{n}{2}}(2\pi|\xi|)}{|\xi|^{\frac{n}{2}}},$$

in view of the result in Appendix B.3. More generally, for $\operatorname{Re} \lambda > -1$, let

$$m_\lambda(\xi) = \begin{cases} (1-|\xi|^2)^\lambda & \text{for } |\xi| \leq 1, \\ 0 & \text{for } |\xi| > 1. \end{cases}$$

Then

$$m_\lambda^\vee(x) = \frac{2\pi}{|x|^{\frac{n-2}{2}}} \int_0^1 J_{\frac{n}{2}-1}(2\pi|x|r)r^{\frac{n}{2}}(1-r^2)^\lambda dr = \frac{\Gamma(\lambda+1)}{\pi^\lambda} \frac{J_{\frac{n}{2}+\lambda}(2\pi|x|)}{|x|^{\frac{n}{2}+\lambda}},$$

using again the identity in Appendix B.3.

B.6 Bessel Functions of Small Arguments

We seek the behavior of $J_\nu(r)$ as $r \rightarrow 0+$. We fix $\nu \in \mathbf{C}$ with $\operatorname{Re} \nu > -\frac{1}{2}$. Then we have the identity

$$J_\nu(r) = \frac{r^\nu}{2^\nu \Gamma(\nu+1)} + S_\nu(r),$$

where

$$S_\nu(r) = \frac{(r/2)^\nu}{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^{+1} (e^{irt} - 1)(1-t^2)^{\nu-\frac{1}{2}} dt$$

and S_ν satisfies

$$|S_\nu(r)| \leq \frac{2^{-\operatorname{Re} \nu} r^{\operatorname{Re} \nu+1}}{(\operatorname{Re} \nu+1) |\Gamma(\nu+\frac{1}{2})| |\Gamma(\frac{1}{2})|}.$$

To prove this estimate we note that

$$\begin{aligned} J_\nu(r) &= \frac{(r/2)^\nu}{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^{+1} (1-t^2)^{\nu-\frac{1}{2}} dt + S_\nu(r) \\ &= \frac{(r/2)^\nu}{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^\pi (\sin^2 \phi)^{\nu-\frac{1}{2}} (\sin \phi) d\phi + S_\nu(r) \\ &= \frac{(r/2)^\nu}{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})} \frac{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})}{\Gamma(\nu+1)} + S_\nu(r), \end{aligned}$$

where we evaluated the last integral using the result in Appendix A.4. Using that $|e^{irt} - 1| \leq r|t|$, we deduce the assertion regarding the size of $|S_\nu(r)|$.

It follows from these facts and the estimate in Appendix A.7 that for $0 < r \leq 1$ and $\text{Re } \nu > -1/2$ we have

$$|J_\nu(r)| \leq C_0(\text{Re } \nu) e^{\max\left(\frac{1}{(\text{Re } \nu + \frac{1}{2})^2}, \frac{1}{\text{Re } \nu + \frac{1}{2}}\right) |\text{Im } \nu|^2} r^{\text{Re } \nu},$$

where C_0 is a constant that depends smoothly on $\text{Re } \nu \in (-1/2, \infty)$.

B.7 Bessel Functions of Large Arguments

For $r > 0$ and complex numbers ν with $\text{Re } \nu > -1/2$ we prove the identity

$$J_\nu(r) = \frac{(r/2)^\nu}{\Gamma(\nu + \frac{1}{2})\Gamma(\frac{1}{2})} \left[ie^{-ir} \int_0^\infty e^{-rt} (t^2 + 2it)^{\nu - \frac{1}{2}} dt - ie^{ir} \int_0^\infty e^{-rt} (t^2 - 2it)^{\nu - \frac{1}{2}} dt \right].$$

Fix $0 < \delta < 1/10 < 10 < R < \infty$. We consider the region $\Omega_{\delta,R}$ in the complex plane whose boundary is the set consisting of the interval $[-1 + \delta, 1 - \delta]$ union a quarter circle centered at 1 of radius δ from $1 - \delta$ to $1 + i\delta$, union the line segments from $1 + i\delta$ to $1 + iR$, from $1 + iR$ to $-1 + iR$, and from $-1 + iR$ to $-1 + i\delta$, union a quarter circle centered at -1 of radius δ from $-1 + i\delta$ to $-1 + \delta$. This is a simply connected region on the interior of which the holomorphic function $(1 - z^2)$ has no zeros. Since $\Omega_{\delta,R}$ is contained in the complement of the negative imaginary axis, there is a holomorphic branch of the logarithm such that $\log(t)$ is real, $\log(-t) = \log|t| + i\pi$, and $\log(it) = \log|t| + i\pi/2$ for $t > 0$. Since the function $\log(1 - z^2)$ is well defined and holomorphic in $\Omega_{\delta,R}$, we may define the holomorphic function

$$(1 - z^2)^{\nu - \frac{1}{2}} = e^{(\nu - \frac{1}{2})\log(1 - z^2)}$$

for $z \in \Omega_{\delta,R}$. Since $e^{irz}(1 - z^2)^{\nu - \frac{1}{2}}$ has no poles in $\Omega_{\delta,R}$, Cauchy's theorem yields

$$i \int_\delta^R e^{ir(1+it)} (t^2 - 2it)^{\nu - \frac{1}{2}} dt + \int_{-1+\delta}^{1-\delta} e^{irt} (1 - t^2)^{\nu - \frac{1}{2}} dt + i \int_R^\delta e^{ir(-1+it)} (t^2 + 2it)^{\nu - \frac{1}{2}} dt + E(\delta, R) = 0,$$

where $E(\delta, R)$ is the sum of the integrals over the two small quarter-circles of radius δ and the line segment from $1 + iR$ to $-1 + iR$. The first two of these integrals are bounded by constants times δ , the latter by a constant times $R^{2\text{Re } \nu - 1} e^{-rR}$; hence $E(\delta, R) \rightarrow 0$ as $\delta \rightarrow 0$ and $R \rightarrow \infty$. We deduce the identity

$$\int_{-1}^{+1} e^{irt} (1 - t^2)^{\nu - \frac{1}{2}} dt = ie^{-ir} \int_0^\infty e^{-rt} (t^2 + 2it)^{\nu - \frac{1}{2}} dt - ie^{ir} \int_0^\infty e^{-rt} (t^2 - 2it)^{\nu - \frac{1}{2}} dt.$$

Estimating the two integrals on the right by putting absolute values inside and multiplying by the missing factor $r^{\nu} 2^{-\nu} (\Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2}))^{-1}$, we obtain

$$|J_{\nu}(r)| \leq 2 \frac{(r/2)^{\operatorname{Re} \nu} e^{\frac{\pi}{2} |\operatorname{Im} \nu|}}{|\Gamma(\nu + \frac{1}{2})| \Gamma(\frac{1}{2})} \int_0^{\infty} e^{-rt} t^{\operatorname{Re} \nu - \frac{1}{2}} (\sqrt{t^2 + 4})^{\operatorname{Re} \nu - \frac{1}{2}} dt,$$

since the absolute value of the argument of $t^2 \pm 2it$ is at most $\pi/2$. When $\operatorname{Re} \nu > 1/2$, we use the inequality $(\sqrt{t^2 + 4})^{\operatorname{Re} \nu - \frac{1}{2}} \leq 2^{\operatorname{Re} \nu - \frac{3}{2}} (t^{\operatorname{Re} \nu - \frac{1}{2}} + 2^{\operatorname{Re} \nu - \frac{1}{2}})$ to get

$$|J_{\nu}(r)| \leq 2 \frac{(r/2)^{\operatorname{Re} \nu} e^{\frac{\pi}{2} |\operatorname{Im} \nu|}}{|\Gamma(\nu + \frac{1}{2})| \Gamma(\frac{1}{2})} 2^{\operatorname{Re} \nu - \frac{3}{2}} \left[\frac{\Gamma(2\operatorname{Re} \nu)}{r^{2\operatorname{Re} \nu}} + 2^{\operatorname{Re} \nu} \frac{\Gamma(\operatorname{Re} \nu + \frac{1}{2})}{r^{\operatorname{Re} \nu + \frac{1}{2}}} \right].$$

When $1/2 \geq \operatorname{Re} \nu > -1/2$ we use that $(\sqrt{t^2 + 4})^{\operatorname{Re} \nu - \frac{1}{2}} \leq 1$ to deduce that

$$|J_{\nu}(r)| \leq 2 \frac{(r/2)^{\operatorname{Re} \nu} e^{\frac{\pi}{2} |\operatorname{Im} \nu|}}{|\Gamma(\nu + \frac{1}{2})| \Gamma(\frac{1}{2})} \frac{\Gamma(\operatorname{Re} \nu + \frac{1}{2})}{r^{\operatorname{Re} \nu + \frac{1}{2}}}.$$

These estimates yield that for $\operatorname{Re} \nu > -1/2$ and $r \geq 1$ we have

$$|J_{\nu}(r)| \leq C_1 (\operatorname{Re} \nu) e^{(\max((\operatorname{Re} \nu + \frac{1}{2})^{-2}, (\operatorname{Re} \nu + \frac{1}{2})^{-1}) + \frac{\pi}{2}) |\operatorname{Im} \nu|^2} r^{-1/2}$$

using the result in Appendix A.7, where C_1 is a constant that depends smoothly on $\operatorname{Re} \nu$ on the interval $(-1/2, \infty)$.

B.8 Asymptotics of Bessel Functions

We obtain asymptotics for $J_{\nu}(r)$ as $r \rightarrow \infty$ whenever $\operatorname{Re} \nu > -1/2$. We have the following identity for $r > 0$:

$$J_{\nu}(r) = \sqrt{\frac{2}{\pi r}} \cos\left(r - \frac{\pi \nu}{2} - \frac{\pi}{4}\right) + R_{\nu}(r),$$

where R_{ν} is given by

$$\begin{aligned} R_{\nu}(r) &= \frac{(2\pi)^{-\frac{1}{2}} r^{\nu}}{\Gamma(\nu + \frac{1}{2})} e^{i(r - \frac{\pi \nu}{2} - \frac{\pi}{4})} \int_0^{\infty} e^{-rt} t^{\nu + \frac{1}{2}} \left[\left(1 + \frac{it}{2}\right)^{\nu - \frac{1}{2}} - 1 \right] \frac{dt}{t} \\ &\quad + \frac{(2\pi)^{-\frac{1}{2}} r^{\nu}}{\Gamma(\nu + \frac{1}{2})} e^{-i(r - \frac{\pi \nu}{2} - \frac{\pi}{4})} \int_0^{\infty} e^{-rt} t^{\nu + \frac{1}{2}} \left[\left(1 - \frac{it}{2}\right)^{\nu - \frac{1}{2}} - 1 \right] \frac{dt}{t} \end{aligned}$$

and satisfies $|R_{\nu}(r)| \leq C_{\nu} r^{-3/2}$ whenever $r \geq 1$.

To see the validity of this identity we write

$$ie^{-ir}(t^2 + 2it)^{v-\frac{1}{2}} = (2t)^{v-\frac{1}{2}} e^{-i(r-\frac{v\pi}{2}-\frac{\pi}{4})} (1 - \frac{it}{2})^{v-\frac{1}{2}},$$

$$-ie^{ir}(t^2 - 2it)^{v-\frac{1}{2}} = (2t)^{v-\frac{1}{2}} e^{i(r-\frac{v\pi}{2}-\frac{\pi}{4})} (1 + \frac{it}{2})^{v-\frac{1}{2}}.$$

Inserting these expressions into the corresponding integrals in the formula proved in Appendix B.7, adding and subtracting 1 from each term $(1 \pm \frac{it}{2})^{v-\frac{1}{2}}$, and multiplying by the missing factor $(r/2)^v/\Gamma(v + \frac{1}{2})\Gamma(\frac{1}{2})$, we obtain the claimed identity

$$J_v(r) = \sqrt{\frac{2}{\pi r}} \cos\left(r - \frac{\pi v}{2} - \frac{\pi}{4}\right) + R_v(r).$$

It remains to estimate $R_v(r)$. We begin by noting that for a, b real with $a > -1$ we have the pair of inequalities

$$|(1 \pm iy)^{a+ib} - 1| \leq 3(|a| + |b|) \left(2^{\frac{a+1}{2}} e^{\frac{\pi}{2}|b|}\right) y \quad \text{when } 0 < y < 1,$$

$$|(1 \pm iy)^{a+ib} - 1| \leq (1 + y^2)^{\frac{a}{2}} e^{\frac{\pi}{2}|b|} + 1 \leq 2 \left(2^{\frac{a+1}{2}} e^{\frac{\pi}{2}|b|}\right) y^a \quad \text{when } 1 \leq y < \infty.$$

The first inequality is proved by splitting into real and imaginary parts and applying the mean value theorem in the real part. Taking $v - \frac{1}{2} = a + ib$, $y = t/2$, and inserting these estimates into the integrals appearing in R_v , we obtain

$$|R_v(r)| \leq \frac{2^{\frac{1}{2}\text{Re } v + \frac{5}{4}} e^{\frac{\pi}{2}|\text{Im } v|} r^{\text{Re } v}}{(2\pi)^{1/2} |\Gamma(v + \frac{1}{2})|} \left[\frac{3\sqrt{2}|v|}{2} \int_0^2 e^{-rt} t^{\text{Re } v + \frac{3}{2}} \frac{dt}{t} + \frac{2\sqrt{2}}{2^{\text{Re } v}} \int_2^\infty e^{-rt} t^{2\text{Re } v} \frac{dt}{t} \right].$$

It follows that for all $r > 0$ we have

$$|R_v(r)| \leq 2 \frac{2^{\frac{1}{2}\text{Re } v} e^{\frac{\pi}{2}|\text{Im } v|}}{|\Gamma(v + \frac{1}{2})|} \left[|v| \frac{\Gamma(\text{Re } v + \frac{3}{2})}{r^{3/2}} + \frac{r^{-\text{Re } v}}{2^{\text{Re } v}} \int_{2r}^\infty e^{-t} t^{2\text{Re } v} \frac{dt}{t} \right]$$

$$\leq 2 \frac{2^{\frac{1}{2}\text{Re } v} e^{\frac{\pi}{2}|\text{Im } v|}}{|\Gamma(v + \frac{1}{2})|} \left[|v| \frac{\Gamma(\text{Re } v + \frac{3}{2})}{r^{3/2}} + \frac{2^{\text{Re } v} \Gamma(2\text{Re } v)}{r^{\text{Re } v} e^r} \right],$$

using that $e^{-t} \leq e^{-t/2} e^{-r}$ for $t \geq 2r$. We conclude that for $r \geq 1$ and $\text{Re } v > -1/2$ we have

$$|R_v(r)| \leq \frac{2^{\frac{1}{2}\text{Re } v} e^{\frac{\pi}{2}|\text{Im } v|}}{\Gamma(\text{Re } v + \frac{1}{2})} e^{\max((\text{Re } v + \frac{1}{2})^{-2}, (\text{Re } v + \frac{1}{2})^{-1})|\text{Im } v|^2}$$

$$\left[|v| \frac{\Gamma(\text{Re } v + \frac{3}{2})}{r^{3/2}} + \frac{2^{\text{Re } v} \Gamma(2\text{Re } v)}{r^{\text{Re } v} e^r} \right]$$

$$\leq C_2(\text{Re } v) e^{\left(\max((\text{Re } v + \frac{1}{2})^{-2}, (\text{Re } v + \frac{1}{2})^{-1}) + \frac{\pi}{2} + 1\right)|\text{Im } v|^2} r^{-3/2},$$

via the result in Appendix A.7, where C_2 is a constant that depends smoothly on $\text{Re } v$ on the interval $(-1/2, \infty)$.

B.9 Bessel Functions of general complex indices

We now discuss how to extend $J_\nu(t)$ to complex values of ν when $t > 0$. Identity (5)

$$J_\nu(t) = \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \frac{1}{\Gamma(j+\nu+1)} \left(\frac{t}{2}\right)^{2j+\nu}$$

of Appendix B.2 expresses $J_\nu(t)$ as a power series when ν is a complex number with $\operatorname{Re} \nu > -1/2$. But this power series converges absolutely for all complex values of ν by the ratio test. Indeed, for j sufficiently large we have

$$\left| \frac{\frac{(-1)^{j+1}}{(j+1)!} \frac{1}{\Gamma(j+1+\nu+1)} \left(\frac{t}{2}\right)^{2j+2+\nu}}{\frac{(-1)^j}{j!} \frac{1}{\Gamma(j+\nu+1)} \left(\frac{t}{2}\right)^{2j+\nu}} \right| = \left| \frac{\Gamma(j+\nu+1) \left(\frac{t}{2}\right)^2}{(j+1)\Gamma(j+\nu+2)} \right| = \left| \frac{\left(\frac{t}{2}\right)^2}{(j+1)(j+\nu+1)} \right|$$

which tends to zero as $j \rightarrow \infty$, thus the ratio test applies.

Therefore the power series defines an entire function of ν . Since this entire function coincides with J_ν on $(-\frac{1}{2}, \infty) \times \mathbf{R}$, by analytic continuation, $J_\nu(t)$ has an entire extension for all $t > 0$.

We obtain an integral representation of $J_z(t)$ when $-1/2 \geq \operatorname{Re} z > -3/2$. Fix $t > 0$. Then we write

$$\begin{aligned} J_z(t) &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} (1-s^2)^z \frac{ds}{\sqrt{1-s^2}} \\ &= \frac{2\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^1 \cos(ts) (1-s^2)^{z-\frac{1}{2}} ds \\ &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^1 \frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} u^{z-\frac{1}{2}} du, \end{aligned}$$

where the last step follows by the change of variables $u = 1 - s^2$. By the mean value theorem, there is a constant c in $(0, 1)$ such that

$$\begin{aligned} &\frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^1 \frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} u^{z-\frac{1}{2}} du \\ &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\sqrt{\pi}} \int_0^1 \frac{d}{du} \left[\frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} \right] (c) u^{z+\frac{1}{2}} du + \frac{\left(\frac{t}{2}\right)^z \cos(t)}{\sqrt{\pi}} \frac{\int_0^1 u^{z-\frac{1}{2}} du}{\Gamma(z+\frac{1}{2})} \\ &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\sqrt{\pi}} \int_0^1 \frac{d}{du} \left[\frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} \right] (c) u^{z+\frac{1}{2}} du + \frac{\left(\frac{t}{2}\right)^z \cos(t)}{\sqrt{\pi}} \frac{1}{\Gamma(z+\frac{3}{2})}, \end{aligned}$$

where the second identity is due to the fact that the integral $\int_0^1 u^{z-\frac{1}{2}} du$ is equal to $(z+\frac{1}{2})^{-1}$ when $\text{Re } z > -1/2$ and thus when divided by $\Gamma(z+\frac{1}{2})$ can be analytically extended to an entire function. It follows from these calculations that

$$J_{-\frac{1}{2}}(t) = \sqrt{\frac{2}{\pi}} \frac{\cos(t)}{\sqrt{t}}.$$

We now estimate $J_z(t)$ when $-3/2 < \text{Re } z \leq -1/2$. Given $q \in (0, 1)$ we write

$$J_z(t) = J_z^1(t; q) + J_z^2(t; q) + J_z^3(t; q),$$

where

$$\begin{aligned} J_z^1(t; q) &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^q \frac{d}{du} \left[\frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} \right] (c) u^{z+\frac{1}{2}} du \\ J_z^2(t; q) &= \frac{\left(\frac{t}{2}\right)^z \cos(t)}{\sqrt{\pi}} \frac{q^{z+\frac{1}{2}}}{\Gamma(z+\frac{3}{2})} \\ J_z^3(t; q) &= \frac{\left(\frac{t}{2}\right)^z}{\Gamma(z+\frac{1}{2})\Gamma(\frac{1}{2})} \int_q^1 \frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} u^{z-\frac{1}{2}} du \end{aligned}$$

for some c satisfying $0 < c < q$.

Suppose that $t > 2$. Then we pick $q = 1/t$. We have

$$|J_z^1(t; 1/t)| \leq \frac{\left(\frac{t}{2}\right)^{\text{Re } z}}{|\Gamma(z+\frac{1}{2})|\sqrt{\pi}} \frac{t}{(\sqrt{1-1/t})^3} \frac{1}{\text{Re } z + \frac{3}{2}} \left(\frac{1}{t}\right)^{\text{Re } z + \frac{3}{2}} \leq \frac{C_1(\text{Re } z) t^{-\frac{1}{2}}}{|\Gamma(z+\frac{1}{2})|},$$

for some constant C_1 depending on $\text{Re } z$. The second term obviously satisfies

$$|J_z^2(t; 1/t)| \leq \frac{C_2(\text{Re } z) t^{-\frac{1}{2}}}{|\Gamma(z+\frac{3}{2})|}.$$

In $J_z^3(t; 1/t)$ we write $\frac{\cos(t\sqrt{1-u})}{\sqrt{1-u}} = \frac{-2}{t} \frac{d}{du} [\sin(t\sqrt{1-u})]$ and integrate by parts to obtain the estimate

$$|J_z^3(t; 1/t)| \leq \frac{\left(\frac{t}{2}\right)^{\text{Re } z}}{|\Gamma(z+\frac{1}{2})|\sqrt{\pi}} \left[\frac{2}{t} + \frac{2}{t^{\frac{1}{2}+\text{Re } z}} \right] \leq \frac{C_3(\text{Re } z) t^{-\frac{1}{2}}}{|\Gamma(z+\frac{1}{2})|}$$

when $-1/2 \geq \text{Re } z > -3/2$. Thus for these z 's and $t > 2$ we have

$$|J_z(t)| \leq C_0(\text{Re } z) t^{-\frac{1}{2}} \left[\frac{1}{|\Gamma(z+\frac{1}{2})|} + \frac{1}{|\Gamma(z+\frac{3}{2})|} \right].$$

When $0 < t \leq 2$ we choose $q = 1/2$. Then each of $|J_z^k(t; 1/2)|$, $k = 1, 2, 3$ is bounded by a multiple of $t^{\operatorname{Re} z}$ and we obtain the estimate

$$|J_z(t)| \leq C_{00}(\operatorname{Re} z) t^{\operatorname{Re} z} \left[\frac{1}{|\Gamma(z + \frac{1}{2})|} + \frac{1}{|\Gamma(z + \frac{3}{2})|} \right].$$

when $-1/2 \geq \operatorname{Re} z > -3/2$ and $0 < t \leq 2$.

In the special case $\operatorname{Re} z = -1/2$ or $z = -1/2 + i\theta$, $\theta \in \mathbf{R}$, we deduce

$$|J_{-\frac{1}{2}+i\theta}(t)| \leq C t^{-\frac{1}{2}} \left[\frac{1}{|\Gamma(1+i\theta)|} + \frac{1}{|\Gamma(i\theta)|} \right] \leq C t^{-\frac{1}{2}} (1 + |\theta|) e^{|\theta|^2}$$

for some constant C independent of all parameters; see Appendix A.7 for the last inequality.

Appendix C

Rademacher Functions

C.1 Definition of the Rademacher Functions

The Rademacher functions are defined on $[0, 1]$ as follows: $r_0(t) = 1$; $r_1(t) = 1$ for $0 \leq t \leq 1/2$ and $r_1(t) = -1$ for $1/2 < t \leq 1$; $r_2(t) = 1$ for $0 \leq t \leq 1/4$, $r_2(t) = -1$ for $1/4 < t \leq 1/2$, $r_2(t) = 1$ for $1/2 < t \leq 3/4$, and $r_2(t) = -1$ for $3/4 < t \leq 1$; and so on. According to this definition, we have that $r_j(t) = \text{sgn}(\sin(2^j \pi t))$ for $j = 0, 1, 2, \dots$. It is easy to check that the r_j 's are mutually independent random variables on $[0, 1]$. This means that for all functions f_j we have

$$\int_0^1 \prod_{j=0}^n f_j(r_j(t)) dt = \prod_{j=0}^n \int_0^1 f_j(r_j(t)) dt.$$

To see the validity of this identity, we write its right-hand side as

$$\begin{aligned} f_0(1) \prod_{j=1}^n \int_0^1 f_j(r_j(t)) dt &= f_0(1) \prod_{j=1}^n \frac{f_j(1) + f_j(-1)}{2} \\ &= \frac{f_0(1)}{2^n} \sum_{S \subset \{1, 2, \dots, n\}} \prod_{j \in S} f_j(1) \prod_{j \notin S} f_j(-1) \end{aligned}$$

and we observe that there is a one-to-one and onto correspondence between subsets S of $\{1, 2, \dots, n\}$ and intervals $I_k = [\frac{k}{2^n}, \frac{k+1}{2^n}]$, $k = 0, 1, \dots, 2^n - 1$, such that the restriction of the function $\prod_{j=1}^n f_j(r_j(t))$ on I_k is equal to

$$\prod_{j \in S} f_j(1) \prod_{j \notin S} f_j(-1).$$

It follows that the last of the three equal displayed expressions is

$$f_0(1) \sum_{k=0}^{2^n-1} \int_{I_k} \prod_{j=1}^n f_j(r_j(t)) dt = \int_0^1 \prod_{j=0}^n f_j(r_j(t)) dt.$$

C.2 Khintchine's Inequalities

The following property of the Rademacher functions is of fundamental importance and with far-reaching consequences in analysis:

For any $0 < p < \infty$ and for any real-valued square summable sequences $\{a_j\}$ and $\{b_j\}$ we have

$$B_p \left(\sum_j |a_j + ib_j|^2 \right)^{\frac{1}{2}} \leq \left\| \sum_j (a_j + ib_j) r_j \right\|_{L^p([0,1])} \leq A_p \left(\sum_j |a_j + ib_j|^2 \right)^{\frac{1}{2}}$$

for some constants $0 < A_p, B_p < \infty$ that depend only on p .

These inequalities reflect the orthogonality of the Rademacher functions in L^p (especially when $p \neq 2$). Khintchine [193] was the first to prove a special form of this inequality, and he used it to estimate the asymptotic behavior of certain random walks. Later this inequality was systematically studied almost simultaneously by Littlewood [226] and by Paley and Zygmund [274], who proved the more general form stated previously. The foregoing inequalities are usually referred to by Khintchine's name.

C.3 Derivation of Khintchine's Inequalities

Both assertions in Appendix C.2 can be derived from an exponentially decaying distributional inequality for the function

$$F(t) = \sum_j (a_j + ib_j) r_j(t), \quad t \in [0, 1],$$

when a_j, b_j are square summable real numbers.

We first obtain a distributional inequality for the above function F under the following three assumptions:

- (a) The sequence $\{b_j\}$ is identically zero.
- (b) All but finitely many terms of the sequence $\{a_j\}$ are zero.
- (c) The sequence $\{a_j\}$ satisfies $(\sum_j |a_j|^2)^{1/2} = 1$.

Let $\rho > 0$. Under assumptions (a), (b), and (c), independence gives

$$\begin{aligned} \int_0^1 e^{\rho \sum_j a_j r_j(t)} dt &= \prod_j \int_0^1 e^{\rho a_j r_j(t)} dt \\ &= \prod_j \frac{e^{\rho a_j} + e^{-\rho a_j}}{2} \\ &\leq \prod_j e^{\frac{1}{2} \rho^2 a_j^2} = e^{\frac{1}{2} \rho^2 \sum_j a_j^2} = e^{\frac{1}{2} \rho^2}, \end{aligned}$$

where we used the inequality $\frac{1}{2}(e^x + e^{-x}) \leq e^{\frac{1}{2}x^2}$ for all real x , which can be checked using power series expansions. Since the same argument is also valid for $-\sum a_j r_j(t)$, we obtain that

$$\int_0^1 e^{\rho|F(t)|} dt \leq 2e^{\frac{1}{2}\rho^2}.$$

From this it follows that

$$e^{\rho\alpha} |\{t \in [0, 1] : |F(t)| > \alpha\}| \leq \int_0^1 e^{\rho|F(t)|} dt \leq 2e^{\frac{1}{2}\rho^2}$$

and hence we obtain the distributional inequality

$$d_F(\alpha) = |\{t \in [0, 1] : |F(t)| > \alpha\}| \leq 2e^{\frac{1}{2}\rho^2 - \rho\alpha} = 2e^{-\frac{1}{2}\alpha^2},$$

by picking $\rho = \alpha$. The L^p norm of F can now be computed easily. Formula (1.1.6) gives

$$\|F\|_{L^p}^p = \int_0^\infty p\alpha^{p-1} d_F(\alpha) d\alpha \leq \int_0^\infty p\alpha^{p-1} 2e^{-\frac{\alpha^2}{2}} d\alpha = 2^{\frac{p}{2}} p \Gamma(p/2).$$

We have now proved that

$$\|F\|_{L^p} \leq \sqrt{2} (p \Gamma(p/2))^{\frac{1}{p}} \|F\|_{L^2}$$

under assumptions (a), (b), and (c).

We now dispose of assumptions (a), (b), and (c). Assumption (b) can be easily eliminated by a limiting argument and (c) by a scaling argument. To dispose of assumption (a), let a_j and b_j be real numbers. We have

$$\begin{aligned} \left\| \sum_j (a_j + ib_j)r_j \right\|_{L^p} &\leq \left\| \left| \sum_j a_j r_j \right| + \left| \sum_j b_j r_j \right| \right\|_{L^p} \\ &\leq \left\| \sum_j a_j r_j \right\|_{L^p} + \left\| \sum_j b_j r_j \right\|_{L^p} \\ &\leq \sqrt{2} (p \Gamma(p/2))^{\frac{1}{p}} \left(\left(\sum_j |a_j|^2 \right)^{\frac{1}{2}} + \left(\sum_j |b_j|^2 \right)^{\frac{1}{2}} \right) \\ &\leq \sqrt{2} (p \Gamma(p/2))^{\frac{1}{p}} \sqrt{2} \left(\sum_j |a_j + ib_j|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Let us now set $A_p = 2(p \Gamma(p/2))^{1/p}$ when $p > 2$. Since we have the trivial estimate $\|F\|_{L^p} \leq \|F\|_{L^2}$ when $0 < p \leq 2$, we obtain the required inequality $\|F\|_{L^p} \leq A_p \|F\|_{L^2}$ with

$$A_p = \begin{cases} 1 & \text{when } 0 < p \leq 2, \\ 2 p^{\frac{1}{p}} \Gamma(p/2)^{\frac{1}{p}} & \text{when } 2 < p < \infty. \end{cases}$$

Using Sterling’s formula in Appendix A.6, we see that A_p is asymptotic to \sqrt{p} as $p \rightarrow \infty$.

We now discuss the converse inequality $B_p \|F\|_{L^2} \leq \|F\|_{L^p}$. It is clear that $\|F\|_{L^2} \leq \|F\|_{L^p}$ when $p \geq 2$ and we may therefore take $B_p = 1$ for $p \geq 2$. Let us now consider the case $0 < p < 2$. Pick an s such that $2 < s < \infty$. Find a $0 < \theta < 1$ such that

$$\frac{1}{2} = \frac{1 - \theta}{p} + \frac{\theta}{s}.$$

Then

$$\|F\|_{L^2} \leq \|F\|_{L^p}^{1-\theta} \|F\|_{L^s}^\theta \leq \|F\|_{L^p}^{1-\theta} A_s^\theta \|F\|_{L^2}^\theta.$$

It follows that

$$\|F\|_{L^2} \leq A_s^{\frac{\theta}{1-\theta}} \|F\|_{L^p}.$$

We have now proved the inequality $B_p \|F\|_{L^2} \leq \|F\|_{L^p}$ with

$$B_p = \begin{cases} 1 & \text{when } 2 \leq p < \infty, \\ \sup_{s>2} A_s^{-\frac{\frac{1}{p}-\frac{1}{2}}{\frac{1}{2}-\frac{1}{s}}} & \text{when } 0 < p < 2. \end{cases}$$

Observe that the function $s \rightarrow A_s^{-\left(\frac{1}{p}-\frac{1}{2}\right)/\left(\frac{1}{2}-\frac{1}{s}\right)}$ tends to zero as $s \rightarrow 2+$ and as $s \rightarrow \infty$. Hence it must attain its maximum for some $s = s(p)$ in the interval $(2, \infty)$. We see that $B_p \geq 16 \cdot 256^{-1/p}$ when $p < 2$ by taking $s = 4$.

It is worthwhile to mention that the best possible values of the constants A_p and B_p in Khintchine’s inequality are known when $b_j = 0$. In this case Szarek [353] showed that the best possible value of B_1 is $1/\sqrt{2}$, and later Haagerup [141] found that when $b_j = 0$ the best possible values of A_p and B_p are the numbers

$$A_p = \begin{cases} 1 & \text{when } 0 < p \leq 2, \\ 2^{\frac{1}{2}} \pi^{-\frac{1}{2p}} \Gamma\left(\frac{p+1}{2}\right) & \text{when } 2 < p < \infty, \end{cases}$$

and

$$B_p = \begin{cases} 2^{\frac{1}{2}-\frac{1}{p}} & \text{when } 0 < p \leq p_0, \\ 2^{\frac{1}{2}} \pi^{-\frac{1}{2p}} \Gamma\left(\frac{p+1}{2}\right) & \text{when } p_0 < p < 2, \\ 1 & \text{when } 2 < p < \infty, \end{cases}$$

where $p_0 = 1.84742\dots$ is the unique solution of the equation $2\Gamma\left(\frac{p+1}{2}\right) = \sqrt{\pi}$ in the interval $(1, 2)$.

C.4 Khintchine’s Inequalities for Weak Type Spaces

We note that the following weak type estimates are valid:

$$4^{-\frac{1}{p}} B_{\frac{p}{2}} \left(\sum_j |a_j + ib_j|^2 \right)^{\frac{1}{2}} \leq \left\| \sum_j (a_j + ib_j) r_j \right\|_{L^{p,\infty}} \leq A_p \left(\sum_j |a_j + ib_j|^2 \right)^{\frac{1}{2}}$$

for all $0 < p < \infty$.

Indeed, the upper estimate is a simple consequence of the fact that L^p is a subspace of $L^{p,\infty}$. For the converse inequality we use the fact that $L^{p,\infty}([0, 1])$ is contained in $L^{p/2}([0, 1])$ and we have (see Exercise 1.1.11)

$$\|F\|_{L^{p/2}} \leq 4^{\frac{1}{p}} \|F\|_{L^{p,\infty}}.$$

Since any Lorentz space $L^{p,q}([0, 1])$ can be sandwiched between $L^{2p}([0, 1])$ and $L^{p/2}([0, 1])$, similar inequalities hold for all Lorentz spaces $L^{p,q}([0, 1])$, $0 < p < \infty$, $0 < q \leq \infty$.

C.5 Extension to Several Variables

We first extend the inequality on the right in Appendix C.2 to several variables. For a positive integer n we let

$$F_n(t_1, \dots, t_n) = \sum_{j_1} \cdots \sum_{j_n} c_{j_1, \dots, j_n} r_{j_1}(t_1) \cdots r_{j_n}(t_n),$$

for $t_j \in [0, 1]$, where c_{j_1, \dots, j_n} is a sequence of complex numbers and F_n is a function defined on $[0, 1]^n$.

For any $0 < p < \infty$ and for any complex-valued square summable sequence of n variables $\{c_{j_1, \dots, j_n}\}_{j_1, \dots, j_n}$, we have the following inequalities for F_n :

$$B_p^n \left(\sum_{j_1} \cdots \sum_{j_n} |c_{j_1, \dots, j_n}|^2 \right)^{\frac{1}{2}} \leq \|F_n\|_{L^p([0,1]^n)} \leq A_p^n \left(\sum_{j_1} \cdots \sum_{j_n} |c_{j_1, \dots, j_n}|^2 \right)^{\frac{1}{2}},$$

where A_p, B_p are the constants in Appendix C.2. The norms are over $[0, 1]^n$.

The case $n = 2$ is indicative of the general case. For $p \geq 2$ we have

$$\begin{aligned} \int_0^1 \int_0^1 |F_2(t_1, t_2)|^p dt_1 dt_2 &\leq A_p^p \int_0^1 \left(\sum_{j_1} \left| \sum_{j_2} c_{j_1, j_2} r_{j_2}(t_2) \right|^2 \right)^{\frac{p}{2}} dt_1 \\ &\leq A_p^p \left(\sum_{j_1} \left(\int_0^1 \left| \sum_{j_2} c_{j_1, j_2} r_{j_2}(t_2) \right|^p dt_2 \right)^{\frac{2}{p}} \right)^{\frac{p}{2}} \\ &\leq A_p^{2p} \left(\sum_{j_1} \sum_{j_2} |c_{j_1, j_2}|^2 \right)^{\frac{p}{2}}, \end{aligned}$$

where we used Minkowski's integral inequality (with exponent $p/2 \geq 1$) in the second inequality and the result in the case $n = 1$ twice.

The case $p < 2$ follows trivially from Hölder's inequality with constant $A_p = 1$. The reverse inequalities follow exactly as in the case of one variable. Replacing A_p by A_p^n in the argument, giving the reverse inequality in the case $n = 1$, we obtain the constant B_p^n .

Likewise one may extend the weak type inequalities of Appendix C.3 in several variables.

Appendix D

Spherical Coordinates

D.1 Spherical Coordinate Formula

Switching integration from spherical coordinates to Cartesian is achieved via the following identity:

$$\int_{R\mathbb{S}^{n-1}} f(x) d\sigma(x) = \int_{\varphi_1=0}^{\pi} \cdots \int_{\varphi_{n-2}=0}^{\pi} \int_{\varphi_{n-1}=0}^{2\pi} f(x(\varphi)) J(n, R, \varphi) d\varphi_{n-1} \cdots d\varphi_1,$$

where

$$\begin{aligned} x_1 &= R \cos \varphi_1, \\ x_2 &= R \sin \varphi_1 \cos \varphi_2, \\ x_3 &= R \sin \varphi_1 \sin \varphi_2 \cos \varphi_3, \\ &\dots \\ x_{n-1} &= R \sin \varphi_1 \sin \varphi_2 \sin \varphi_3 \cdots \sin \varphi_{n-2} \cos \varphi_{n-1}, \\ x_n &= R \sin \varphi_1 \sin \varphi_2 \sin \varphi_3 \cdots \sin \varphi_{n-2} \sin \varphi_{n-1}, \end{aligned}$$

and $0 \leq \varphi_1, \dots, \varphi_{n-2} \leq \pi, 0 \leq \varphi_{n-1} = \theta < 2\pi,$

$$x(\varphi) = (x_1(\varphi_1, \dots, \varphi_{n-1}), \dots, x_n(\varphi_1, \dots, \varphi_{n-1})),$$

and

$$\begin{aligned} J(n, R, \varphi) &= \left(\sum_{k=1}^n \left| \det \left(\frac{\partial x_i}{\partial \varphi_j} \right)_{\substack{1 \leq i \neq k \leq n \\ 1 \leq j \leq n-1}} \right|^2 \right)^{\frac{1}{2}} \\ &= R^{n-1} (\sin \varphi_1)^{n-2} \cdots (\sin \varphi_{n-3})^2 (\sin \varphi_{n-2}) \end{aligned}$$

is the Jacobian of the transformation.

D.2 A Useful Change of Variables Formula

The following formula is useful in computing integrals over the sphere \mathbf{S}^{n-1} when $n \geq 2$. Let f be a function defined on \mathbf{S}^{n-1} . Then we have

$$\int_{R\mathbf{S}^{n-1}} f(x) d\sigma(x) = \int_{-R}^{+R} \int_{\sqrt{R^2-s^2}\mathbf{S}^{n-2}} f(s, \theta) d\theta \frac{R ds}{\sqrt{R^2-s^2}}.$$

To prove this formula, let $\varphi' = (\varphi_2, \dots, \varphi_{n-1})$ and

$$x' = x'(\varphi') = (\cos \varphi_2, \sin \varphi_2 \cos \varphi_3, \dots, \sin \varphi_2 \cdots \sin \varphi_{n-2} \sin \varphi_{n-1}).$$

Using the change of variables in Appendix D.1 we express

$$\int_{R\mathbf{S}^{n-1}} f(x) d\sigma(x)$$

as the iterated integral

$$\int_{\varphi_1=0}^{\pi} \left[\int_{\varphi_2=0}^{\pi} \cdots \int_{\varphi_{n-1}=0}^{2\pi} f(R \cos \varphi_1, R \sin \varphi_1 x'(\varphi')) J(n-1, 1, \varphi') d\varphi' \right] \frac{R d\varphi_1}{(R \sin \varphi_1)^{2-n}},$$

and we can realize the expression inside the square brackets as

$$\int_{\mathbf{S}^{n-2}} f(R \cos \varphi_1, R \sin \varphi_1 x') d\sigma(x').$$

Consequently,

$$\int_{R\mathbf{S}^{n-1}} f(x) d\sigma(x) = \int_{\varphi_1=0}^{\pi} \int_{\mathbf{S}^{n-2}} f(R \cos \varphi_1, R \sin \varphi_1 x') d\sigma(x') R^{n-1} (\sin \varphi_1)^{n-2} d\varphi_1,$$

and the change of variables

$$s = R \cos \varphi_1, \quad \varphi_1 \in (0, \pi),$$

$$ds = -R \sin \varphi_1 d\varphi_1, \quad \sqrt{R^2 - s^2} = R \sin \varphi_1,$$

yields

$$\int_{R\mathbf{S}^{n-1}} f(x) d\sigma(x) = \int_{-R}^R \left\{ \int_{\mathbf{S}^{n-2}} f(s, \sqrt{R^2 - s^2} \theta) d\theta \right\} (\sqrt{R^2 - s^2})^{n-2} \frac{R ds}{\sqrt{R^2 - s^2}}.$$

Rescaling the sphere \mathbf{S}^{n-2} to $\sqrt{R^2 - s^2} \mathbf{S}^{n-2}$ yields the claimed identity.

D.3 Computation of an Integral over the Sphere

Let K be a function on the line. We use the result in Appendix D.2 to show that for $n \geq 2$ we have

$$\int_{\mathbf{S}^{n-1}} K(x \cdot \theta) d\theta = \frac{2\pi^{\frac{n-1}{2}}}{\Gamma(\frac{n-1}{2})} \int_{-1}^{+1} K(s|x|)(\sqrt{1-s^2})^{n-3} ds$$

when $x \in \mathbf{R}^n \setminus \{0\}$. Let $x' = x/|x|$ and pick a matrix $A \in O(n)$ such that $Ae_1 = x'$, where $e_1 = (1, 0, \dots, 0)$. We have

$$\begin{aligned} \int_{\mathbf{S}^{n-1}} K(x \cdot \theta) d\theta &= \int_{\mathbf{S}^{n-1}} K(|x|(x' \cdot \theta)) d\theta \\ &= \int_{\mathbf{S}^{n-1}} K(|x|(Ae_1 \cdot \theta)) d\theta \\ &= \int_{\mathbf{S}^{n-1}} K(|x|(e_1 \cdot A^{-1}\theta)) d\theta \\ &= \int_{\mathbf{S}^{n-1}} K(|x|\theta_1) d\theta \\ &= \int_{-1}^{+1} K(|x|s) \omega_{n-2} (\sqrt{1-s^2})^{n-2} \frac{ds}{\sqrt{1-s^2}} \\ &= \omega_{n-2} \int_{-1}^{+1} K(s|x|) (\sqrt{1-s^2})^{n-3} ds, \end{aligned}$$

where $\omega_{n-2} = 2\pi^{\frac{n-1}{2}} \Gamma(\frac{n-1}{2})^{-1}$ is the surface area of \mathbf{S}^{n-2} .

For example, we have

$$\int_{\mathbf{S}^{n-1}} \frac{d\theta}{|\xi \cdot \theta|^\alpha} = \omega_{n-2} \int_{-1}^{+1} \frac{1}{|s|^\alpha |\xi|^\alpha} (1-s^2)^{\frac{n-3}{2}} ds = \frac{1}{|\xi|^\alpha} \frac{2\pi^{\frac{n-1}{2}} \Gamma(\frac{1-\alpha}{2})}{\Gamma(\frac{n-\alpha}{2})},$$

and the integral converges only when $\operatorname{Re} \alpha < 1$.

D.4 The Computation of Another Integral over the Sphere

We compute the following integral for $n \geq 2$:

$$\int_{\mathbf{S}^{n-1}} \frac{d\theta}{|\theta - e_1|^\alpha},$$

where $e_1 = (1, 0, \dots, 0)$. Applying the formula in Appendix D.2, we obtain

$$\begin{aligned} \int_{\mathbf{S}^{n-1}} \frac{d\theta}{|\theta - e_1|^\alpha} &= \int_{-1}^{+1} \int_{\theta \in \sqrt{1-s^2} \mathbf{S}^{n-2}} \frac{d\theta}{(|s-1|^2 + |\theta|^2)^{\frac{\alpha}{2}}} \frac{ds}{\sqrt{1-s^2}} \\ &= \int_{-1}^{+1} \omega_{n-2} \frac{(1-s^2)^{\frac{n-2}{2}}}{((1-s)^2 + 1-s^2)^{\frac{\alpha}{2}}} \frac{ds}{\sqrt{1-s^2}} \\ &= \frac{\omega_{n-2}}{2^{\frac{\alpha}{2}}} \int_{-1}^{+1} \frac{(1-s^2)^{\frac{n-3}{2}}}{(1-s)^{\frac{\alpha}{2}}} ds \\ &= \frac{\omega_{n-2}}{2^{\frac{\alpha}{2}}} \int_{-1}^{+1} (1-s)^{\frac{n-3-\alpha}{2}} (1+s)^{\frac{n-3}{2}} ds, \end{aligned}$$

which converges exactly when $\operatorname{Re} \alpha < n-1$.

D.5 Integration over a General Surface

Suppose that S is a hypersurface in \mathbf{R}^n of the form $S = \{(u, \Phi(u)) : u \in D\}$, where D is an open subset of \mathbf{R}^{n-1} and Φ is a continuously differentiable mapping from D to \mathbf{R} . Let σ be the canonical surface measure on S . If g is a function on S , then we have

$$\int_S g(y) d\sigma(y) = \int_D g(x, \Phi(x)) \left(1 + \sum_{j=1}^{n-1} |\partial_j \Phi(x)|^2\right)^{\frac{1}{2}} dx.$$

Specializing to the sphere, we obtain

$$\int_{\mathbf{S}^{n-1}} g(\theta) d\theta = \int_{\substack{\xi' \in \mathbf{R}^{n-1} \\ |\xi'| < 1}} \left[g(\xi', \sqrt{1-|\xi'|^2}) + g(\xi', -\sqrt{1-|\xi'|^2}) \right] \frac{d\xi'}{\sqrt{1-|\xi'|^2}}.$$

D.6 The Stereographic Projection

Define a map $\Pi : \mathbf{R}^n \rightarrow \mathbf{S}^n$ by the formula

$$\Pi(x_1, \dots, x_n) = \left(\frac{2x_1}{1+|x|^2}, \dots, \frac{2x_n}{1+|x|^2}, \frac{|x|^2-1}{1+|x|^2} \right).$$

It is easy to see that Π is a one-to-one map from \mathbf{R}^n onto the sphere \mathbf{S}^n minus the north pole $e_{n+1} = (0, \dots, 0, 1)$. Its inverse is given by the formula

$$\Pi^{-1}(\theta_1, \dots, \theta_{n+1}) = \left(\frac{\theta_1}{1-\theta_{n+1}}, \dots, \frac{\theta_n}{1-\theta_{n+1}} \right).$$

The Jacobian of the map is verified to be $J_{\Pi}(x) = \left(\frac{2}{1+|x|^2}\right)^{-n}$, and the following change of variables formulas are valid:

$$\int_{\mathbf{S}^n} F(\theta) d\theta = \int_{\mathbf{R}^n} F(\Pi(x)) J_{\Pi}(x) dx$$

and

$$\int_{\mathbf{R}^n} F(x) dx = \int_{\mathbf{S}^n} F(\Pi^{-1}(\theta)) J_{\Pi^{-1}}(\theta) d\theta,$$

where

$$J_{\Pi^{-1}}(\theta) = \frac{1}{J_{\Pi}(\Pi^{-1}(\theta))} = \left(\frac{|\theta_1|^2 + \cdots + |\theta_n|^2 + |1 - \theta_{n+1}|^2}{2|1 - \theta_{n+1}|^2} \right)^n.$$

Another interesting formula about the stereographic projection Π is

$$|\Pi(x) - \Pi(y)| = 2|x - y|(1 + |x|^2)^{-1/2}(1 + |y|^2)^{-1/2}, \quad x, y \in \mathbf{R}^n.$$

Appendix E

Some Trigonometric Identities and Inequalities

The following inequalities are valid for t real:

$$\begin{aligned}
 0 < t < \frac{\pi}{2} &\implies \sin(t) < t < \tan(t), \\
 0 < |t| < \frac{\pi}{2} &\implies \frac{2}{\pi} < \frac{\sin(t)}{t} < 1, \\
 -\infty < t < +\infty &\implies |\sin(t)| \leq |t|, \\
 -\infty < t < +\infty &\implies |1 - \cos(t)| \leq \frac{|t|^2}{2}, \\
 -\infty < t < +\infty &\implies |1 - e^{it}| \leq |t|, \\
 |t| \leq \frac{\pi}{2} &\implies |\sin(t)| \geq \frac{2|t|}{\pi}, \\
 |t| \leq \pi &\implies |1 - \cos(t)| \geq \frac{2|t|^2}{\pi^2}, \\
 |t| \leq \pi &\implies |1 - e^{it}| \geq \frac{2|t|}{\pi}.
 \end{aligned}$$

The following sum to product formulas are valid:

$$\begin{aligned}
 \sin(a) + \sin(b) &= 2 \sin\left(\frac{a+b}{2}\right) \cos\left(\frac{a-b}{2}\right), \\
 \sin(a) - \sin(b) &= 2 \cos\left(\frac{a+b}{2}\right) \sin\left(\frac{a-b}{2}\right), \\
 \cos(a) + \cos(b) &= 2 \cos\left(\frac{a+b}{2}\right) \cos\left(\frac{a-b}{2}\right), \\
 \cos(a) - \cos(b) &= -2 \sin\left(\frac{a+b}{2}\right) \sin\left(\frac{a-b}{2}\right).
 \end{aligned}$$

The following identities are also easily proved:

$$\sum_{k=1}^N \cos(kx) = -\frac{1}{2} + \frac{\sin((N + \frac{1}{2})x)}{2 \sin(\frac{x}{2})},$$
$$\sum_{k=1}^N \sin(kx) = \frac{\cos(\frac{x}{2}) - \cos((N + \frac{1}{2})x)}{2 \sin(\frac{x}{2})}.$$

Appendix F

Summation by Parts

Let $\{a_k\}_{k=0}^{\infty}$, $\{b_k\}_{k=0}^{\infty}$ be two sequences of complex numbers. Then for $N \geq 1$ we have

$$\sum_{k=0}^N a_k b_k = A_N b_N - \sum_{k=0}^{N-1} A_k (b_{k+1} - b_k),$$

where

$$A_k = \sum_{j=0}^k a_j.$$

More generally we have

$$\sum_{k=M}^N a_k b_k = A_N b_N - A_{M-1} b_M - \sum_{k=M}^{N-1} A_k (b_{k+1} - b_k),$$

whenever $0 \leq M \leq N$, where $A_{-1} = 0$ and

$$A_k = \sum_{j=0}^k a_j$$

for $k \geq 0$.

Appendix G

Basic Functional Analysis

A quasi-norm is a nonnegative functional $\|\cdot\|$ on a vector space X that satisfies $\|x+y\|_X \leq K(\|x\|_X + \|y\|_X)$ for some $K \geq 0$ and all $x, y \in X$ and also $\|\lambda x\|_X = |\lambda| \|x\|_X$ for all scalars λ . When $K = 1$, the quasi-norm is called a norm. A quasi-Banach space is a quasi-normed space that is complete with respect to the topology generated by the quasi-norm. The proofs of the following theorems can be found in several books including Albiac and Kalton [1], Kalton Peck and Roberts [188], and Rudin [306].

The Hahn–Banach theorem. Let X be a normed vector space (over the real or complex numbers), let Y be a subspace of X , and let P be a positively homogeneous subadditive¹ functional on X . Then for every linear functional Λ on Y that satisfies

$$|\Lambda(y)| \leq P(y)$$

for all $y \in Y$, there is a linear functional Λ' on X such that

$$\begin{aligned} \Lambda'(y) &= \Lambda(y) && \text{for all } y \in Y, \\ |\Lambda'(x)| &\leq P(x) && \text{for all } x \in X. \end{aligned}$$

In particular, every bounded linear functional on a subspace has an extension on the entire space with the same norm.

Banach–Alaoglu theorem. Let X be a normed vector space and let X^* be the space of all bounded linear functionals on X . Then the closed unit ball of X^* is compact in the weak* topology.

A special case is the sequential version of this theorem, which asserts that the closed unit ball of the dual space of a separable normed vector space is sequentially compact in the weak* topology. Indeed, the weak* topology on the closed unit ball

¹ this means that $P(x) \geq 0$ for all $x \in X$, $P(\lambda x) = \lambda P(x)$ for all $\lambda > 0$ and all $x \in X$, and that $P(x+z) \leq P(x) + P(z)$ for all $x, z \in X$.

of the dual of a separable space is metrizable, and thus compactness and sequential compactness are equivalent.

Open mapping theorem. Suppose that X and Y are quasi-Banach spaces and T is a bounded surjective linear map from X onto Y . Then T maps open sets to open sets, i.e., it is an open mapping. Moreover, if T is injective, then there exists a constant $K < \infty$ such that for all $x \in X$ we have

$$\|x\|_X \leq K\|T(x)\|_Y.$$

Closed graph theorem. Suppose that X and Y are quasi-Banach spaces and T is a linear map from X to Y whose graph is a closed set, i.e., whenever $x_k, x \in X$ and $(x_k, T(x_k)) \mapsto (x, y)$ in $X \times Y$ for some $y \in Y$, then $T(x) = y$. Then T is a bounded linear map from X to Y .

Uniform boundedness principle. Suppose that X is a quasi-Banach space, Y is a quasi-normed space and $(T_\alpha)_{\alpha \in I}$ is a family of bounded linear maps from X to Y such that for all $x \in X$ there exists a $C_x < \infty$ such that

$$\sup_{\alpha \in I} \|T_\alpha(x)\|_Y \leq C_x.$$

Then there exists a constant $K < \infty$ such that

$$\sup_{\alpha \in I} \|T_\alpha\|_{X \rightarrow Y} \leq K.$$

Appendix H

The Minimax Lemma

Minimax type results are used in the theory of games and have their origin in the work of Von Neumann [369]. Much of the theory in this subject is based on convex analysis techniques. For instance, this is the case with the next proposition, which is needed in the “difficult” inequality in the proof of the minimax lemma. We refer to Fan [111] for a general account of minimax results. The following exposition is based on the simple presentation in Appendix A2 of [122].

Minimax Lemma. *Let A, B be convex subsets of certain vector spaces. Assume that a topology is defined in B for which it is a compact Hausdorff space and assume that there is a function $\Phi : A \times B \rightarrow \mathbf{R} \cup \{+\infty\}$ that satisfies the following:*

- (a) $\Phi(\cdot, b)$ is a concave function on A for each $b \in B$,
- (b) $\Phi(a, \cdot)$ is a convex function on B for each $a \in A$,
- (c) $\Phi(a, \cdot)$ is lower semicontinuous on B for each $a \in A$.

Then the following identity holds:

$$\min_{b \in B} \sup_{a \in A} \Phi(a, b) = \sup_{a \in A} \min_{b \in B} \Phi(a, b).$$

To prove the lemma we need the following proposition:

Proposition. *Let B be a convex compact subset of a vector space and suppose that $g_j : B \rightarrow \mathbf{R} \cup \{+\infty\}$, $j = 1, 2, \dots, n$, are convex and lower semicontinuous functions. If*

$$\max_{1 \leq j \leq n} g_j(b) > 0 \quad \text{for all } b \in B,$$

then there exist nonnegative numbers $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$$\lambda_1 g_1(b) + \lambda_2 g_2(b) + \dots + \lambda_n g_n(b) > 0 \quad \text{for all } b \in B.$$

Proof. We first consider the case $n = 2$. Define subsets of B

$$B_1 = \{b \in B : g_1(b) \leq 0\}, \quad B_2 = \{b \in B : g_2(b) \leq 0\}.$$

If $B_1 = \emptyset$, we take $\lambda_1 = 1$ and $\lambda_2 = 0$, and we similarly deal with the case $B_2 = \emptyset$. If B_1 and B_2 are nonempty, then they are closed and thus compact. The hypothesis of the proposition implies that $g_2(b) > 0 \geq g_1(b)$ for all $b \in B_1$. Therefore, the function $-g_1(b)/g_2(b)$ is well defined and upper semicontinuous on B_1 and thus attains its maximum. The same is true for $-g_2(b)/g_1(b)$ defined on B_2 . We set

$$\mu_1 = \max_{b \in B_1} \frac{-g_1(b)}{g_2(b)} \geq 0, \quad \mu_2 = \max_{b \in B_2} \frac{-g_2(b)}{g_1(b)} \geq 0.$$

We need to find $\lambda > 0$ such that $\lambda g_1(b) + g_2(b) > 0$ for all $b \in B$. This is clearly satisfied if $b \notin B_1 \cup B_2$, while for $b_1 \in B_1$ and $b_2 \in B_2$ we have

$$\begin{aligned} \lambda g_1(b_1) + g_2(b_1) &\geq (1 - \lambda \mu_1) g_2(b_1), \\ \lambda g_1(b_2) + g_2(b_2) &\geq (\lambda - \mu_2) g_1(b_2). \end{aligned}$$

Therefore, it suffices to find a $\lambda > 0$ such that $1 - \lambda \mu_1 > 0$ and $\lambda - \mu_2 > 0$. Such a λ exists if and only if $\mu_1 \mu_2 < 1$. To prove that $\mu_1 \mu_2 < 1$, we can assume that $\mu_1 \neq 0$ and $\mu_2 \neq 0$. Then we take $b_1 \in B_1$ and $b_2 \in B_2$, for which the maxima μ_1 and μ_2 are attained, respectively. Then we have

$$\begin{aligned} g_1(b_1) + \mu_1 g_2(b_1) &= 0, \\ g_1(b_2) + \frac{1}{\mu_2} g_2(b_2) &= 0. \end{aligned}$$

But $g_1(b_1) < 0 < g_1(b_2)$; thus taking $b_\theta = \theta b_1 + (1 - \theta) b_2$ for some θ in $(0, 1)$, we have

$$g_1(b_\theta) \leq \theta g_1(b_1) + (1 - \theta) g_1(b_2) = 0.$$

Considering the same convex combination of the last displayed equations and using this identity, we obtain that

$$\mu_1 \mu_2 \theta g_2(b_1) + (1 - \theta) g_2(b_2) = 0.$$

The hypothesis of the proposition implies that $g_2(b_\theta) > 0$ and the convexity of g_2 :

$$\theta g_2(b_1) + (1 - \theta) g_2(b_2) > 0.$$

Since $g_2(b_1) > 0$, we must have $\mu_1 \mu_2 g_2(b_1) < g_2(b_1)$, which gives $\mu_1 \mu_2 < 1$. This proves the required claim and completes the case $n = 2$.

We now use induction to prove the proposition for arbitrary n . Assume that the result has been proved for $n - 1$ functions. Consider the subset of B

$$B_n = \{b \in B : g_n(b) \leq 0\}.$$

If $B_n = \emptyset$, we choose $\lambda_1 = \lambda_2 = \dots = \lambda_{n-1} = 0$ and $\lambda_n = 1$. If B_n is not empty, then it is compact and convex and we can restrict g_1, g_2, \dots, g_{n-1} to B_n . Using the induction hypothesis, we can find $\lambda_1, \lambda_2, \dots, \lambda_{n-1} \geq 0$ such that

$$g_0(b) = \lambda_1 g_1(b) + \lambda_2 g_2(b) + \cdots + \lambda_{n-1} g_{n-1}(b) > 0$$

for all $b \in B_n$. Then g_0 and g_n are convex lower semicontinuous functions on B , and $\max(g_0(b), g_n(b)) > 0$ for all $b \in B$. Using the case $n = 2$, which was first proved, we can find $\lambda_0, \lambda_n \geq 0$ such that for all $b \in B$ we have

$$\begin{aligned} 0 &< \lambda_0 g_0(b) + \lambda_n g_n(b) \\ &= \lambda_0 \lambda_1 g_1(b) + \lambda_0 \lambda_2 g_2(b) + \cdots + \lambda_0 \lambda_{n-1} g_{n-1}(b) + \lambda_n g_n(b). \end{aligned}$$

This establishes the case of n functions and concludes the proof of the induction and hence of the proposition. □

We now turn to the proof of the minimax lemma.

Proof. The fact that the left-hand side in the required conclusion of the minimax lemma is at least as big as the right-hand side is obvious. We can therefore concentrate on the converse inequality. In doing this we may assume that the right-hand side is finite. Without loss of generality we can subtract a finite constant from $\Phi(a, b)$, and so we can also assume that

$$\sup_{a \in A} \min_{b \in B} \Phi(a, b) = 0.$$

Then, by hypothesis (c) of the minimax lemma, the subsets

$$B_a = \{b \in B : \Phi(a, b) \leq 0\}, \quad a \in A$$

of B are closed and nonempty, and we show that they satisfy the finite intersection property. Indeed, suppose that

$$B_{a_1} \cap B_{a_2} \cap \cdots \cap B_{a_n} = \emptyset$$

for some $a_1, a_2, \dots, a_n \in A$. We write $g_j(b) = \Phi(a_j, b)$, $j = 1, 2, \dots, n$, and we observe that the conditions of the previous proposition are satisfied. Therefore we can find $\lambda_1, \lambda_2, \dots, \lambda_n \geq 0$ such that for all $b \in B$ we have

$$\lambda_1 \Phi(a_1, b) + \lambda_2 \Phi(a_2, b) + \cdots + \lambda_n \Phi(a_n, b) > 0.$$

For simplicity we normalize the λ_j 's by setting $\lambda_1 + \lambda_2 + \cdots + \lambda_n = 1$. If we set $a_0 = \lambda_1 a_1 + \lambda_2 a_2 + \cdots + \lambda_n a_n$, the concavity hypothesis (a) gives

$$\Phi(a_0, b) > 0$$

for all $b \in B$, contradicting the fact that $\sup_{a \in A} \min_{b \in B} \Phi(a, b) = 0$. Therefore, the family of closed subsets $\{B_a\}_{a \in A}$ of B satisfies the finite intersection property. The compactness of B now implies $\bigcap_{a \in A} B_a \neq \emptyset$. Take $b_0 \in \bigcap_{a \in A} B_a$. Then $\Phi(a, b_0) \leq 0$ for every $a \in A$, and therefore

$$\min_{b \in B} \sup_{a \in A} \Phi(a, b) \leq \sup_{a \in A} \Phi(a, b_0) \leq 0$$

as required. □

Appendix I

Taylor's and Mean Value Theorem in Several Variables

I.1 Multivariable Taylor's Theorem

For a multiindex $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbf{Z}^+ \cup \{0\})^n$, we denote by $|\alpha| = \alpha_1 + \dots + \alpha_n$ its size, we define $\alpha! = \alpha_1! \cdots \alpha_n!$ its factorial, and we set

$$h^\alpha = h_1^{\alpha_1} \cdots h_n^{\alpha_n},$$

where $h = (h_1, \dots, h_n)$; here $0^0 = 1$.

Let $k \in \mathbf{Z}^+ \cup \{0\}$. Suppose a real-valued \mathcal{C}^{k+1} function f is defined on an open convex subset Ω of \mathbf{R}^n . Suppose that $x \in \Omega$ and $x + h \in \Omega$. Then we have the *Taylor expansion formula*

$$f(x+h) = \sum_{|\alpha| \leq k} \frac{\partial^\alpha f(x)}{\alpha!} h^\alpha + R(h, x, k),$$

where the remainder $R(h, x, k)$ can be expressed either in Lagrange's mean value form

$$R(h, x, k) = \sum_{|\alpha|=k+1} \frac{\partial^\alpha f(x+ch)}{\alpha!} h^\alpha$$

for some $c \in (0, 1)$, or in integral form

$$R(h, x, k) = (k+1) \sum_{|\alpha|=k+1} \frac{h^\alpha}{\alpha!} \int_0^1 (1-t)^k \partial^\alpha f(x+th) dt.$$

I.2 The Mean value Theorem

Suppose that f is as above and $k = 0$. Then for given $x, y \in \Omega$ we have

$$f(y) - f(x) = \int_0^1 \nabla f((1-t)x + ty) \cdot (y-x) dt = \nabla f((1-c)x + cy) \cdot (y-x)$$

for some $c \in (0, 1)$. This is a special case of Taylor's formula when $k = 0$.

Appendix J

The Whitney Decomposition of Open Sets in \mathbf{R}^n

J.1 Decomposition of Open Sets

An arbitrary open set in \mathbf{R}^n can be decomposed as a union of disjoint cubes whose lengths are proportional to their distance from the boundary of the open set. See, for instance, Figure J.1 when the open set is the unit disk in \mathbf{R}^2 . For a given cube Q in \mathbf{R}^n , we denote by $\ell(Q)$ its length.

Proposition. *Let Ω be an open nonempty proper subset of \mathbf{R}^n . Then there exists a family of closed dyadic cubes $\{Q_j\}_j$ (called the Whitney cubes of Ω) such that*

- (a) $\bigcup_j Q_j = \Omega$ and the Q_j 's have disjoint interiors.
- (b) $\sqrt{n}\ell(Q_j) \leq \text{dist}(Q_j, \Omega^c) \leq 4\sqrt{n}\ell(Q_j)$. Thus $10\sqrt{n}Q_j$ meets Ω^c .
- (c) If the boundaries of two cubes Q_j and Q_k touch, then

$$\frac{1}{4} \leq \frac{\ell(Q_j)}{\ell(Q_k)} \leq 4.$$

- (d) For a given Q_j there exist at most $12^n - 4^n$ cubes Q_k that touch it.
- (e) Let $0 < \varepsilon < 1/4$. If Q_j^* has the same center as Q_j and $\ell(Q_j^*) = (1 + \varepsilon)\ell(Q_j)$ then

$$\chi_\Omega = \sum_j \chi_{Q_j^*} \leq 12^n - 4^n + 1.$$

Proof. Let \mathcal{D}_k be the collection of all dyadic cubes of the form

$$\{(x_1, \dots, x_n) \in \mathbf{R}^n : m_j 2^{-k} \leq x_j < (m_j + 1)2^{-k}\},$$

where $m_j \in \mathbf{Z}$. Observe that each cube in \mathcal{D}_k gives rise to 2^n cubes in \mathcal{D}_{k+1} by bisecting each side.

Write the set Ω as the union of the sets

$$\Omega_k = \{x \in \Omega : 2\sqrt{n}2^{-k} < \text{dist}(x, \Omega^c) \leq 4\sqrt{n}2^{-k}\}$$

over all $k \in \mathbf{Z}$. Let \mathcal{F}' be the set of all cubes Q in \mathcal{D}_k for some $k \in \mathbf{Z}$ such that $Q \cap \Omega_k \neq \emptyset$. We show that the collection \mathcal{F}' satisfies property (b). Let $Q \in \mathcal{F}'$ and pick $x \in \Omega_k \cap Q$ for some $k \in \mathbf{Z}$. Observe that

$$\sqrt{n}2^{-k} \leq \text{dist}(x, \Omega^c) - \sqrt{n} \ell(Q) \leq \text{dist}(Q, \Omega^c) \leq \text{dist}(x, \Omega^c) \leq 4\sqrt{n}2^{-k},$$

which proves (b).

Next we observe that

$$\bigcup_{Q \in \mathcal{F}'} Q = \Omega.$$

Indeed, every Q in \mathcal{F}' is contained in Ω (since it has positive distance from its complement) and every $x \in \Omega$ lies in some Ω_k and in some dyadic cube in \mathcal{D}_k .




Fig. J.1 The Whitney decomposition of the unit disk.

The problem is that the cubes in the collection \mathcal{F}' may not be disjoint. We have to refine the collection \mathcal{F}' by eliminating those cubes that are contained in some other cubes in the collection. Recall that two dyadic cubes have disjoint interiors or else one contains the other. For every cube Q in \mathcal{F}' we can therefore consider the unique *maximal* cube Q^{\max} in \mathcal{F}' that contains it. Two different such maximal cubes must have disjoint interiors by maximality. Now set $\mathcal{F} = \{Q^{\max} : Q \in \mathcal{F}'\}$.

The collection of cubes $\{Q_j\}_{j \in \mathcal{F}}$ clearly satisfies (a) and (b), and we now turn our attention to the proof of (c). Observe that if Q_j and Q_k in \mathcal{F} touch then

$$\sqrt{n} \ell(Q_j) \leq \text{dist}(Q_j, \Omega^c) \leq \text{dist}(Q_j, Q_k) + \text{dist}(Q_k, \Omega^c) \leq 0 + 4\sqrt{n} \ell(Q_k),$$

which proves (c). To prove (d), note that any cube Q in \mathcal{D}_k is touched by exactly $3^n - 1$ other cubes in \mathcal{D}_k . But each cube Q in \mathcal{D}_k can contain at most 4^n cubes of \mathcal{F} of length at least one-quarter of the length of Q . This fact combined with (c) yields (d). To prove (e), notice that each Q_j^* is contained in Ω by part (b). If $x \in \Omega$, then

$x \in Q_{k_0}$ for some k_0 . If Q_j does not touch Q_{k_0} , then Q_j^* does not touch Q_{k_0} as well. Consequently, the given $x \in \Omega$ may lie only in Q_k^* for these cubes Q_k that touch Q_{k_0} and there are $12^n - 4^n + 1$ such cubes including Q_{k_0} . \square

J.2 Partition of Unity adapted to Whitney cubes

Let us fix an ε such that $0 < \varepsilon < 1/4$. For each cube Q we denote by Q^* the cube with the same center as Q and side length $(1 + \varepsilon)\ell(Q)$, where $\ell(Q)$ is the side length of Q . Let us fix a nonnegative smooth function ϕ that is equal to 1 on the unit cube $Q_0 = [-1/2, 1/2]^n$ and equal to zero outside Q_0^* .

Let $\{Q_k\}_k$ be the family of Whitney cubes of Ω . We denote by c_k the center of Q_k and by ℓ_k its side length. Since $\text{dist}(Q_k, \Omega^c) \geq \sqrt{n}\ell_k$, and $\text{dist}((Q_k^*)^c, Q_k) \leq \varepsilon\sqrt{n}\ell_k$, it follows that

$$\text{dist}(Q_k^*, \Omega^c) \geq (1 - \varepsilon)\sqrt{n}\ell_k > 0,$$

hence Q_k^* is contained in Ω . Since the union of Q_k is Ω , then the union of Q_k^* is also Ω .

For each k we define

$$\phi_k(x) = \phi\left(\frac{x - c_k}{\ell_k}\right)$$

and we set

$$\Phi(x) = \sum_k \phi_k(x).$$

We notice that Φ is smooth and that $\Phi \geq 1$ on Ω . Then we define

$$\varphi_k(x) = \frac{\phi_k(x)}{\Phi(x)}.$$

Obviously φ_k are supported in Q_k^* , the union of Q_k^* is Ω , and we have

$$\sum_k \varphi_k = \chi_\Omega.$$

We would like to control $|\partial^\alpha \varphi_k(x)|$ in terms of ℓ_k . To achieve this we use the Leibniz rule of differentiation. For a fixed k we have that

$$\frac{\partial^\alpha}{\partial x^\alpha} \varphi_k(x) = \sum_{\beta \leq \alpha} \binom{\alpha_1}{\beta_1} \cdots \binom{\alpha_n}{\beta_n} \left(\frac{\partial^\beta}{\partial x^\beta} \frac{1}{\Phi(x)} \right) \frac{\partial^{\alpha-\beta}}{\partial x^{\alpha-\beta}} \phi_k(x)$$

and obviously $|\partial^{\alpha-\beta} \phi_k(x)| \leq \ell_k^{-|\alpha|+|\beta|}$. A simple inductive argument shows that

$$\left| \frac{\partial^\beta}{\partial x^\beta} \frac{1}{\Phi(x)} \right| \leq C \sum_{\substack{0 \leq q_j \leq |\beta| \\ q_1 + \cdots + q_n = |\beta|}} \frac{|\partial_1^{q_1} \cdots \partial_n^{q_n} \Phi(x)|}{\Phi(x)^2},$$

for some constant C , and since for a given $x \in \Omega$, $\Phi(x)$ is the sum of at most 12^n functions with nonzero values, it follows that

$$|\partial_1^{q_1} \dots \partial_n^{q_n} \Phi(x)| \leq C_n \ell_k^{-(q_1 + \dots + q_n)}$$

when $x \in Q_k^*$ and thus

$$\left| \frac{\partial^\beta}{\partial x^\beta} \frac{1}{\Phi(x)} \right| \leq C'_{n,\beta} \ell_k^{-|\beta|}$$

for $x \in Q_k^*$. We conclude that for every multiindex α there is a constant $C_{\alpha,n}$ such that

$$\left| \frac{\partial^\alpha}{\partial x^\alpha} \varphi_k(x) \right| \leq C_{\alpha,n} \ell^{-|\alpha|}.$$

More on Whitney decompositions can be found in the article of Whitney [373] and the books of Stein [338], Krantz and Parks [204].

Glossary

$A \subseteq B$	A is a subset of B (also denoted by $A \subset B$)
$A \subsetneq B$	A is a proper subset of B
$A \supset B$	B is a proper subset of A
A^c	the complement of a set A
χ_E	the characteristic function of the set E
d_f	the distribution function of a function f
f^*	the decreasing rearrangement of a function f
$f_n \uparrow f$	f_n increases monotonically to a function f
\mathbf{Z}	the set of all integers
\mathbf{Z}^+	the set of all positive integers $\{1, 2, 3, \dots\}$
\mathbf{Z}^n	the n -fold product of the integers
\mathbf{R}	the set of real numbers
\mathbf{R}^+	the set of positive real numbers
\mathbf{R}^n	the Euclidean n -space
\mathbf{Q}	the set of rationals
\mathbf{Q}^n	the set of n -tuples with rational coordinates
\mathbf{C}	the set of complex numbers
\mathbf{C}^n	the n -fold product of complex numbers
\mathbf{T}	the unit circle identified with the interval $[0, 1]$
\mathbf{T}^n	the n -dimensional torus $[0, 1]^n$
$ x $	$\sqrt{ x_1 ^2 + \dots + x_n ^2}$ when $x = (x_1, \dots, x_n) \in \mathbf{R}^n$

\mathbf{S}^{n-1}	the unit sphere $\{x \in \mathbf{R}^n : x = 1\}$
e_j	the vector $(0, \dots, 0, 1, 0, \dots, 0)$ with 1 in the j th entry and 0 elsewhere
$\log t$	the logarithm with base e of $t > 0$
$\log_a t$	the logarithm with base a of $t > 0$ ($1 \neq a > 0$)
$\log^+ t$	$\max(0, \log t)$ for $t > 0$
$[t]$	the integer part of the real number t
$x \cdot y$	the quantity $\sum_{j=1}^n x_j y_j$ when $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$
$B(x, R)$	the ball of radius R centered at x in \mathbf{R}^n
ω_{n-1}	the surface area of the unit sphere \mathbf{S}^{n-1}
v_n	the volume of the unit ball $\{x \in \mathbf{R}^n : x < 1\}$
$ A $	the Lebesgue measure of the set $A \subseteq \mathbf{R}^n$
dx	Lebesgue measure
$\text{Avg}_B f$	the average $\frac{1}{ B } \int_B f(x) dx$ of f over the set B
$\langle f, g \rangle$	the real inner product $\int_{\mathbf{R}^n} f(x)g(x) dx$
$\langle f g \rangle$	the complex inner product $\int_{\mathbf{R}^n} f(x)\overline{g(x)} dx$
$\langle u, f \rangle$	the action of a distribution u on a function f
p'	the number $p/(p-1)$, whenever $0 < p \neq 1 < \infty$
$1'$	the number ∞
∞'	the number 1
$f = O(g)$	means $ f(x) \leq M g(x) $ for some M for x near x_0
$f = o(g)$	means $ f(x) / g(x) ^{-1} \rightarrow 0$ as $x \rightarrow x_0$
A^t	the transpose of the matrix A
A^*	the conjugate transpose of a complex matrix A
A^{-1}	the inverse of the matrix A
$O(n)$	the space of real matrices satisfying $A^{-1} = A^t$
$\ T\ _{X \rightarrow Y}$	the norm of the (bounded) operator $T : X \rightarrow Y$
$A \approx B$	means that there exists a $c > 0$ such that $c^{-1} \leq \frac{B}{A} \leq c$
$ \alpha $	indicates the size $ \alpha_1 + \dots + \alpha_n $ of a multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$
$\partial_j^m f$	the m th partial derivative of $f(x_1, \dots, x_n)$ with respect to x_j
$\partial^\alpha f$	$\partial_1^{\alpha_1} \dots \partial_n^{\alpha_n} f$

\mathcal{C}^k	the space of functions f with $\partial^\alpha f$ continuous for all $ \alpha \leq k$
\mathcal{C}_0	the space of continuous functions with compact support
\mathcal{C}_{00}	the space of continuous functions that vanish at infinity
\mathcal{C}_0^∞	the space of smooth functions with compact support
\mathcal{D}	the space of smooth functions with compact support
\mathcal{S}	the space of Schwartz functions
\mathcal{S}_0	the space of Schwartz functions φ with the property $\int_{\mathbf{R}^n} x^\gamma \varphi(x) dx = 0$ for all multi-indices γ .
\mathcal{C}^∞	the space of smooth functions $\bigcup_{k=1}^\infty \mathcal{C}^k$
$\mathcal{D}'(\mathbf{R}^n)$	the space of distributions on \mathbf{R}^n
$\mathcal{S}'(\mathbf{R}^n)$	the space of tempered distributions on \mathbf{R}^n
$\mathcal{E}'(\mathbf{R}^n)$	the space of distributions with compact support on \mathbf{R}^n
\mathcal{P}	the set of all complex-valued polynomials of n real variables
$\mathcal{S}'(\mathbf{R}^n)/\mathcal{P}$	the space of tempered distributions on \mathbf{R}^n modulo polynomials
$\ell(Q)$	the side length of a cube Q in \mathbf{R}^n
∂Q	the boundary of a cube Q in \mathbf{R}^n
$L^p(X, \mu)$	the Lebesgue space over the measure space (X, μ)
$L^p(\mathbf{R}^n)$	the space $L^p(\mathbf{R}^n, \cdot)$
$L^{p,q}(X, \mu)$	the Lorentz space over the measure space (X, μ)
$L^p_{\text{loc}}(\mathbf{R}^n)$	the space of functions that lie in $L^p(K)$ for any compact set K in \mathbf{R}^n
$ d\mu $	the total variation of a finite Borel measure μ on \mathbf{R}^n
$\mathcal{M}(\mathbf{R}^n)$	the space of all finite Borel measures on \mathbf{R}^n
$\mathcal{M}_p(\mathbf{R}^n)$	the space of L^p Fourier multipliers, $1 \leq p \leq \infty$
$\mathcal{M}^{p,q}(\mathbf{R}^n)$	the space of translation-invariant operators that map $L^p(\mathbf{R}^n)$ to $L^q(\mathbf{R}^n)$
$\ \mu\ _{\mathcal{M}}$	$\int_{\mathbf{R}^n} d\mu $ the norm of a finite Borel measure μ on \mathbf{R}^n
\mathcal{M}	the centered Hardy–Littlewood maximal operator with respect to balls
M	the uncentered Hardy–Littlewood maximal operator with respect to balls
\mathcal{M}_c	the centered Hardy–Littlewood maximal operator with respect to cubes
M_c	the uncentered Hardy–Littlewood maximal operator with respect to cubes

\mathcal{M}_μ	the centered maximal operator with respect to a measure μ
M_μ	the uncentered maximal operator with respect to a measure μ
M_s	the strong maximal operator
M_d	the dyadic maximal operator

References

1. Albiac, F., Kalton, N. J., *Topics in Banach Space Theory*, Graduate Texts in Mathematics, 233, Springer, New York, 2006.
2. Aldaz, J. M., *The weak type $(1,1)$ bounds for the maximal function associated to cubes grow to infinity with the dimension*, Ann. of Math. (2nd Ser.) **173** (2011), no. 2, 1013–1023.
3. Alimov, Sh. A., Ashurov, R. R., Pulatov, A. K., *Multiple Fourier series and Fourier integrals*, Proc. Commutative Harmonic Analysis, IV, pp. 1–95, Encyclopaedia Math. Sci. 42, Springer, Berlin, 1992.
4. Alvarez, J., Pérez, C., *Estimates with A_∞ weights for various singular integral operators*, Boll. Un. Mat. Ital. A (7) **8** (1994), no. 1, 123–133.
5. Antonov, N. Yu., *Convergence of Fourier series*, Proceedings of the XX Workshop on Function Theory (Moscow, 1995), East J. Approx. **2** (1996), no. 2, pp. 187–196.
6. Antonov, N. Yu., *The behavior of partial sums of trigonometric Fourier series* [Russian], Ph. D. Thesis, Institute of Mathematics and Mechanics, Ural Branch of the Russian Academy of Sciences, Ekaterinburg, 1998.
7. Aoki, T., *Locally bounded spaces*, Proc. Imp. Acad. Tokyo **18** (1942), no. 10, 585–634.
8. Arhipov, G. I., Karachuba, A. A., Čubarikov, V. N., *Trigonometric integrals*, Math. USSR Izvestija **15** (1980), 211–239.
9. Arias de Reyna, J., *Pointwise convergence of Fourier series*, J. London Math. Soc. (2) **65** (2002), no. 1, 139–153.
10. Arias de Reyna, J., *Pointwise Convergence of Fourier Series*, Lecture Notes in Mathematics, 1785, Springer-Verlag, Berlin, 2002.
11. Ash, J. M., *Multiple trigonometric series*, Studies in Harmonic Analysis (Proc. Conf., DePaul Univ., Chicago, IL, 1974), pp. 76–96, MAA Stud. Math., Vol. 13, Math. Assoc. Amer., Washington, D. C., 1976.
12. Asmar, N., Berkson, E., Gillespie, T. A., *Summability methods for transferring Fourier multipliers and transference of maximal inequalities*, Analysis and Partial Differential Equations, pp. 1–34, Lecture Notes in Pure and Appl. Math. 122, Dekker, New York, 1990.
13. Asmar, N., Berkson, E., Gillespie, T. A., *On Jodeit's multiplier extension theorems*, J. Anal. Math. **64** (1994), 337–345.
14. Astala, K., Iwaniec, T., Saksman, E., *Beltrami operators in the plane*, Duke Math. J. **107** (2001), no. 1, 27–56.
15. Aubrun, G., *Maximal inequality for high-dimensional cubes*, Confluentes Math. **1** (2009), no. 2, 169–179.
16. Babenko, K. I., *An inequality in the theory of Fourier integrals* [Russian], Izv. Akad. Nauk SSSR Ser. Mat. **25** (1961), 531–542.
17. Baernstein II, A., *Some sharp inequalities for conjugate functions*, Indiana Univ. Math. J. **27** (1978), 833–852.

18. Bagby, R., Kurtz, D. S., *Covering lemmas and the sharp function*, Proc. Amer. Math. Soc. **93** (1985), no. 2, 291–296.
19. Bagby, R., Kurtz, D. S., *A rearranged good- λ inequality*, Trans. Amer. Math. Soc. **293** (1986), no. 1, 71–81.
20. Bary, N., *A Treatise on Trigonometric Series*, Vols. I, II, Authorized translation by Margaret F. Mullins, A Pergamon Press Book, The Macmillan Co., New York, 1964.
21. Beckner, W., *Inequalities in Fourier analysis*, Ann. of Math. (2nd Ser.) **102** (1975), no. 1, 159–182.
22. Benedek, A., Calderón, A.-P., Panzone, R., *Convolution operators on Banach-space valued functions*, Proc. Nat. Acad. Sci. U.S.A. **48** (1962), 356–365.
23. Benedetto, J., Frazier, M. (eds.), *Wavelets: Mathematics and Applications*, CRC Press, Boca Raton, FL, 1994.
24. Bennett, C., Sharpley, R., *Interpolation of Operators*, Pure and Applied Mathematics, 129 Academic Press, Inc., Boston, MA, 1988.
25. Bergh, J., Löfström, J., *Interpolation Spaces, An Introduction*, Grundlehren der Mathematischen Wissenschaften, No. 223, Springer-Verlag, Berlin-New York, 1976.
26. Bernstein, S., *Sur la convergence absolue des séries trigonométriques*, Comptes Rendus des Séances de l' Académie des Sciences, Paris, **158** (1914), 1661–1663.
27. Besicovitch, A., *A general form of the covering principle and relative differentiation of additive functions*, Proc. of Cambridge Philos. Soc. **41** (1945), 103–110.
28. Boas, R. P., Bochner, S., *On a theorem of M. Riesz for Fourier series*, J. London Math. Soc. **14** (1939), no. 1, 62–73.
29. Bôcher, M., *Introduction to the theory of Fourier's series*, Ann. of Math. (2nd Ser.) **7** (1906), no. 3, 81–152.
30. Bochner, S., *Summation of multiple Fourier series by spherical means*, Trans. Amer. Math. Soc. **40** (1936), no. 2, 175–207.
31. Bochner, S., *Harmonic Analysis and the Theory of Probability*, University of California Press, Berkeley and Los Angeles, 1955.
32. Bochner, S., *Lectures on Fourier Integrals, (with an author's supplement on monotonic functions, Stieltjes integrals, and harmonic analysis)*, Translated by Morris Tenenbaum and Harry Pollard. Annals of Mathematics Studies, No. 42, Princeton University Press, Princeton, NJ, 1959.
33. Bojarski, B., *Remarks on Markov's inequalities and some properties of polynomials* (Russian summary), Bull. Polish Acad. Sci. Math. **33** (1985), no. 7–8, 355–365.
34. Bourgain, J., *On square functions on the trigonometric system*, Bull. Soc. Math. Belg. Sér. B **37** (1985), no. 1, 20–26.
35. Bourgain, J., *Estimations de certaines fonctions maximales*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 10, 499–502.
36. Bourgain, J., *Averages in the plane over convex curves and maximal operators*, J. Analyse Math. **47** (1986), 69–85.
37. Bourgain, J., *On the Hardy-Littlewood maximal function for the cube*, Israel J. Math. **203** (2014), 275–293.
38. Buckley, S. M., *Estimates for operator norms on weighted spaces and reverse Jensen inequalities*, Trans. Amer. Math. Soc. **340** (1993), no. 1, 253–272.
39. Burkholder, D. L., *Martingale transforms*, Ann. Math. Statist. **37** (1966), 1494–1505.
40. Burkholder, D. L., Gundy, R. F., *Extrapolation and interpolation of quasilinear operators on martingales*, Acta Math. **124** (1970), no. 1, 249–304.
41. Calderón, A. P., *On the theorems of M. Riesz and Zygmund*, Proc. Amer. Math. Soc. **1** (1950), 533–535.
42. Calderón, A. P., *Intermediate spaces and interpolation, the complex method*, Studia Math. **24** (1964), 113–190.
43. Calderón, A. P., *Singular integrals*, Bull. Amer. Math. Soc. **72** (1966), 427–465.
44. Calderón, A. P., *Algebras of singular integral operators*, Singular Integrals (Proc. Sympos. Pure Math. (Chicago, IL, 1966), pp. 18–55, Amer. Math. Soc., Providence, RI, 1967.

45. Calderón, A. P., Torchinsky, A., *Parabolic maximal functions associated with a distribution, II*, *Advances in Math.* **24** (1977), no. 2, 101–171.
46. Calderón, A. P., Zygmund, A., *On the existence of certain singular integrals*, *Acta Math.* **88** (1952), no. 1, 85–139.
47. Calderón, A. P., Zygmund, A., *A note on interpolation of sublinear operators*, *Amer. J. Math.* **78** (1956), 282–288.
48. Calderón, A. P., Zygmund, A., *On singular integrals*, *Amer. J. Math.* **78** (1956), 289–309.
49. Calderón, A. P., Zygmund, A., *Algebras of certain singular integral operators*, *Amer. J. Math.* **78** (1956), 310–320.
50. Calderón, C., *Lacunary spherical means*, *Illinois J. Math.* **23** (1979), no. 3, 476–484.
51. Carbery, A., *An almost-orthogonality principle with applications to maximal functions associated to convex bodies*, *Bull. Amer. Math. Soc. (N.S.)* **14** (1986), no. 2, 269–273.
52. Carbery, A., *Variants of the Calderón–Zygmund theory for L^p -spaces*, *Rev. Mat. Iberoamericana* **2** (1986), no. 4, 381–396.
53. Carbery, A., Christ, M., Wright, J., *Multidimensional van der Corput and sublevel set estimates*, *J. Amer. Math. Soc.* **12** (1999), no. 4, 981–1015.
54. Carleson, L., *On convergence and growth of partial sums of Fourier series*, *Acta Math.* **116** (1966), no. 1, 135–157.
55. Carleson, L., *On the Littlewood–Paley Theorem*, Mittag-Leffler Institute Report, Djursholm, Sweden, 1967.
56. Carro, M. J., *New extrapolation estimates*, *J. Funct. Anal.* **174** (2000), no. 1, 155–166.
57. Carro, M. J., Raposo, J. A., Soria, J., *Recent developments in the theory of Lorentz spaces and weighted inequalities*, *Mem. Amer. Math. Soc.* **187** (2007), no. 877.
58. Carro, M. J., Torres, R. H., Soria, J., *Rubio de Francia’s extrapolation theory: estimates for the distribution function*, *J. Lond. Math. Soc. (2)* **85** (2012), no. 2, 430–454.
59. Christ, M., *A convolution inequality for Cantor–Lebesgue measures*, *Rev. Mat. Iberoamericana* **1** (1985), no. 4, 79–83.
60. Christ, M., *Weak type $(1,1)$ bounds for rough operators I*, *Ann. of Math. (2nd Ser.)* **128** (1988), no. 1, 19–42.
61. Christ, M., Fefferman, R., *A note on weighted norm inequalities for the Hardy–Littlewood maximal operator*, *Proc. Amer. Math. Soc.* **87** (1983), no. 3, 447–448.
62. Christ, M., Grafakos, L., *Best constants for two nonconvolution inequalities*, *Proc. Amer. Math. Soc.* **123** (1995), no. 6, 1687–1693.
63. Christ, M., Rubio de Francia, J. L., *Weak type $(1,1)$ bounds for rough operators II*, *Invent. Math.* **93** (1988), no. 1, 225–237.
64. Chui, C. (ed.), *Wavelets: A Tutorial in Theory and Applications*, Academic Press, San Diego, CA, 1992.
65. Coifman, R. R., *Distribution function inequalities for singular integrals*, *Proc. Nat. Acad. Sci. U.S.A.* **69** (1972), 2838–2839.
66. Coifman, R. R., Fefferman, C., *Weighted norm inequalities for maximal functions and singular integrals*, *Studia Math.* **51** (1974), 241–250.
67. Coifman, R. R., Jones, P., Rubio de Francia, J. L., *Constructive decomposition of BMO functions and factorization of A_p weights*, *Proc. Amer. Math. Soc.* **87** (1983), no. 4, 675–676.
68. Coifman, R. R., Rochberg, R., *Another characterization of BMO*, *Proc. Amer. Math. Soc.* **79** (1980), no. 2, 249–254.
69. Coifman, R. R., Rubio de Francia, J.-L., Semmes, S., *Multiplicateurs de Fourier de $L^p(\mathbb{R})$ et estimations quadratiques*, *C. R. Acad. Sci. Paris Sér. I Math.* **306** (1988), no. 8, 351–354.
70. Coifman, R. R., Weiss, G., *Transference Methods in Analysis*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 31, American Mathematical Society, Providence, RI, 1976.
71. Coifman, R. R., Weiss, G., *Book review of “Littlewood–Paley and multiplier theory”*, *Bull. Amer. Math. Soc.* **84** (1978), no. 2, 242–250.
72. Connett, W. C., *Singular integrals near L^1* , *Harmonic Analysis in Euclidean Spaces*, (Williams Coll., Williamstown, MA, 1978), Part 1, pp. 163–165, *Proc. Sympos. Pure Math.*, XXXV, Part, Amer. Math. Soc., Providence, RI, 1979.

73. Córdoba, A., Fefferman, R., *A geometric proof of the strong maximal theorem*, Ann. of Math. (2nd Ser.) **102** (1975), no. 1, 95–100.
74. Córdoba, A., Fefferman, C., *A weighted norm inequality for singular integrals*, Studia Math. **57** (1976), no. 1, 97–101.
75. Cotlar, M., *A unified theory of Hilbert transforms and ergodic theorems*, Rev. Mat. Cuyana, **1** (1955), 105–167.
76. Cowling, M., Mauceri, G., *On maximal functions*, Rend. Sem. Mat. Fis. Milano **49** (1979), 79–87.
77. Cruz-Uribe, D., *New proofs of two-weight norm inequalities for the maximal operator*, Georgian Math. J. **7** (2000), no. 1, 33–42.
78. Cruz-Uribe, D., Martell, J. M., Pérez, C., *Extrapolation results for A_∞ weights and applications*, J. Funct. Anal. **213** (2004), no. 2, 412–439.
79. Cruz-Uribe, D., Martell, J. M., Pérez, C., *Sharp weighted estimates for approximating dyadic operators*, Electron. Res. Announc. Math. Sci. **17** (2010), 12–19.
80. Cruz-Uribe, D., Martell, J. M., Pérez, C., *Weights, extrapolation and the theory of Rubio de Francia*, Operator Theory: Advances and Applications, 215, Birkhäuser/Springer Basel AG, Basel, 2011.
81. Cruz-Uribe, D., Martell, J. M., Pérez, C., *Sharp weighted estimates for classical operators*, Adv. Math. **229** (2012), no. 1, 408–441.
82. Cruz-Uribe, D., Neugebauer, C. J., *The structure of the reverse Hölder classes*, Trans. Amer. Math. Soc. **347** (1995), no. 8, 2941–2960.
83. Cwikel, M., *The dual of weak L^p* , Ann. Inst. Fourier (Grenoble) **25** (1975), no. 2, 81–126.
84. Cwikel, M., Fefferman, C., *Maximal seminorms on weak L^1* , Studia Math. **69** (1980/81), no. 2, 149–154.
85. Cwikel, M., Fefferman, C., *The canonical seminorm on Weak L^1* , Studia Math. **78** (1984), no. 3, 275–278.
86. Daly, J. E., *A necessary condition for Calderón–Zygmund singular integral operators*, J. Fourier Anal. Appl. **5** (1999), no. 4, 303–308.
87. Daly, J. E., Phillips, K., *On the classification of homogeneous multipliers bounded on $H^1(\mathbb{R}^2)$* , Proc. Amer. Math. Soc. **106** (1989), no. 3, 685–696.
88. Daubechies, I., *Orthonormal bases of compactly supported wavelets*, Comm. Pure Appl. Math. **41** (1988), no. 7, 909–996.
89. Daubechies, I., *Ten Lectures on Wavelets*, CBMS-NSF Regional Conference Series in Applied Math, 61, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1992.
90. Davis, B., *On the integrability of the martingale square function*, Israel J. Math. **8** (1970), 187–190.
91. Davis, B., *On the weak type $(1,1)$ inequality for conjugate functions*, Proc. Amer. Math. Soc. **44** (1974), 307–311.
92. de Guzmán, M., *Real Variable Methods in Fourier Analysis*, North-Holland Mathematics Studies, 46, Notas de Matemática (75), North-Holland Publishing Co., Amsterdam-New York, 1981.
93. de Guzmán, M., *Differentiation of Integrals in \mathbb{R}^n* , Lecture Notes in Math. 481, Springer-Verlag, Berlin-New York, 1975.
94. de Leeuw, K., *On L_p multipliers*, Ann. of Math. (2nd Ser.) **81** (1965), no. 2, 364–379.
95. Diestel, J., Uhl, J. J. Jr., *Vector Measures*, With a foreword by B. J. Pettis, Mathematical Surveys, No. 15, American Mathematical Society, Providence, RI, 1977.
96. Dirichlet, P. G., *Sur la convergence des séries trigonométriques qui servent à représenter une fonction arbitraire entre des limites données*, J. Reine und Angew. Math. **4** (1829), 157–169.
97. Doob, J. L., *Stochastic Processes*, John Wiley & Sons, Inc., New York; Chapman & Hall, Limited, London, 1953.
98. Dragičević, O., Grafakos, L., C. Pereyra, and S. Petermichl, *Extrapolation and sharp norm estimates for classical operators on weighted Lebesgue spaces*, Publ. Mat. **49** (2005), no. 1, 73–91.

99. Dunford, N., Schwartz, J. T., *Linear Operators, I*, General Theory, With the assistance of W. G. Bade and R. G. Bartle, Pure and Applied Mathematics, Vol. 7, Interscience Publishers, Inc., New York; Interscience Publishers, Ltd., London, 1958.
100. Duoandikoetxea, J., *Fourier Analysis*, Grad. Studies in Math. 29, American Mathematical Society, Providence, RI, 2001.
101. Duoandikoetxea, J., *Extrapolation of weights revisited: new proofs and sharp bounds*, J. Funct. Anal. **260** (2011), no. 6, 1886–1901.
102. Duoandikoetxea, J., Rubio de Francia, J.-L., *Maximal and singular integral operators via Fourier transform estimates*, Invent. Math. **84** (1986), no. 3, 541–561.
103. Duoandikoetxea, J., Vega, L., *Spherical means and weighted inequalities*, J. London Math. Soc. (2) **53** (1996), no. 2, 343–353.
104. Duong, X. T., McIntosh, A., *Singular integral operators with non-smooth kernels on irregular domains*, Rev. Mat. Iberoamericana **15** (1999), no. 2, 233–265.
105. Dym, H., McKean, H. P., *Fourier Series and Integrals*, Probability and Mathematical Statistics, No. 14, Academic Press, New York-London, 1972.
106. Edwards, R. E., *Fourier Series: A Modern Introduction. Vol. 1*, 2nd edition, Graduate Texts in Mathematics, 64, Springer-Verlag, New York-Berlin, 1979.
107. Erdélyi, A., *Asymptotic Expansions*, Dover Publications Inc., New York, 1956.
108. Essén, M., *A superharmonic proof of the M. Riesz conjugate function theorem*, Arkiv Math. **22** (1984), no. 2, 241–249.
109. Fan, D., Guo, K., Pan, Y., *A note of a rough singular integral operator*, Math. Inequal. and Appl. **2** (1999), no. 1, 73–81.
110. Fan, D., Pan, Y., *Singular integral operators with rough kernels supported by subvarieties*, Amer J. Math. **119** (1997), no. 4, 799–839.
111. Fan, K., *Minimax theorems*, Proc. Nat. Acad. Sci. U. S. A. **39** (1953), 42–47.
112. Fefferman, C., *On the convergence of multiple Fourier series*, Bull. Amer. Math. Soc. **77** (1971), 744–745.
113. Fefferman, C., *The multiplier problem for the ball*, Ann. of Math. (2nd Ser.) **94** (1971), no. 2, 330–336.
114. Fefferman, C., *The uncertainty principle*, Bull. Amer. Math. Soc. (N. S.) **9** (1983), no. 2, 129–206.
115. Fefferman, C., Stein, E. M., *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107–115.
116. Fefferman, R., *A theory of entropy in Fourier analysis*, Adv. in Math. **30** (1978), no. 3, 171–201.
117. Fefferman, R., *Strong differentiation with respect to measures*, Amer J. Math. **103** (1981), no. 1, 33–40.
118. Folland, G. B., Sitaram, A., *The uncertainty principle: a mathematical survey*, J. Fourier Anal. Appl. **3** (1997), no. 3, 207–238.
119. Fourier, J., *Théorie Analytique de la Chaleur*, Institut de France, Paris, 1822.
120. Frazier, M., *An Introduction to Wavelets Through Linear Algebra*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 1999.
121. García-Cuerva, J., *An extrapolation theorem in the theory of A_p weights*, Proc. Amer. Math. Soc. **87** (1983), no. 3, 422–426.
122. García-Cuerva, J., Rubio de Francia, J.-L., *Weighted Norm Inequalities and Related Topics*, North-Holland Mathematics Studies, 116, Notas de Matemática (104), North-Holland Publishing Co., Amsterdam, 1985.
123. Garnett, J., Jones, P., *The distance in BMO to L^∞* , Ann. of Math. **108** (1978), no. 2, 373–393.
124. Garrigós, G., Seeger, A., *Characterizations of Hankel multipliers*, Math. Ann. **342** (2008), no. 1, 31–68.
125. Gehring, F. W., *The L^p -integrability of the partial derivatives of a quasiconformal mapping*, Acta Math. **130** (1973), no. 1, 265–277.
126. Gelfand, I. M., Šilov, G. E., *Generalized Functions, Vol. 1: Properties and Operations*, Academic Press, New York-London, 1964.
127. Gelfand, I. M., Šilov, G. E., *Generalized Functions, Vol. 2: Spaces of Fundamental and Generalized Functions*, Academic Press, New York-London, 1968.

128. Gibbs, J. W., *Fourier's Series*, Nature **59** (1899), 606.
129. Gohberg, I., Krupnik, N., *Norm of the Hilbert transformation in the L_p space*, Funct. Anal. Appl. **2** (1968), 180–181.
130. Grafakos, L., *Best bounds for the Hilbert transform on $L^p(\mathbb{R}^1)$* , Math. Res. Lett. **4** (1997), no. 4, 469–471.
131. Grafakos, L., *Modern Fourier Analysis*, 3rd edition, Graduate Texts in Math. 250, Springer, New York, 2014.
132. Grafakos, L., Honzík, P., Ryabogin, D., *On the p -independence property of Calderón–Zygmund theory*, J. Reine Angew. Math. **602** (2007), 227–234.
133. Grafakos, L., Kalton, N., *The Marcinkiewicz multiplier condition for bilinear operators*, Studia Math. **146** (2001), no. 2, 115–156.
134. Grafakos, L., Kinnunen, J., *Sharp inequalities for maximal functions associated to general measures*, Proc. Roy. Soc. Edinburgh Sect. A **128** (1998), no. 4, 717–723.
135. Grafakos, L., Martell, J. M., *Extrapolation of weighted norm inequalities for multivariable operators and applications* J. Geom. Anal. **14** (2004), no. 1, 19–46.
136. Grafakos, L., Montgomery-Smith, S., *Best constants for uncentred maximal functions*, Bull. London Math. Soc. **29** (1997), no. 1, 60–64.
137. Grafakos, L., Morpurgo, C., *A Selberg integral formula and applications*, Pacific J. Math. **191** (1999), no. 1, 85–94.
138. Grafakos, L., Stefanov, A., *L^p bounds for singular integrals and maximal singular integrals with rough kernels*, Indiana Univ. Math. J. **47** (1998), no. 2, 455–469.
139. Grafakos, L., Stefanov, A., *Convolution Calderón–Zygmund singular integral operators with rough kernels*, Analysis of Divergence (Orono, ME, 1997), pp. 119–143, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 1999.
140. Gröchenig, K., *Foundations of Time-Frequency Analysis*, Applied and Numerical Harmonic Analysis Birkhäuser Boston, Inc., Boston, MA, 2001.
141. Haagerup, U., *The best constants in Khintchine's inequality*, Studia Math. **70** (1981), no. 3, 231–283.
142. Hahn, L.-S., *On multipliers of p -integrable functions*, Trans. Amer. Math. Soc. **128** (1967), 321–335.
143. Hardy, G. H., *Theorems relating to the summability and convergence of slowly oscillating series*, Proc. London Math. Soc. (2) **8** (1910), no. 1, 301–320.
144. Hardy, G. H., *Note on a theorem of Hilbert*, Math. Z. **6** (1920), no. 3–4, 314–317.
145. Hardy, G. H., *Note on some points in the integral calculus, LX. An inequality between integrals*, Messenger of Math. **54** (1925), 150–156.
146. Hardy, G. H., Littlewood, J. E., *A maximal theorem with function-theoretic applications*, Acta Math. **54** (1930), no. 1, 81–116.
147. Hardy, G. H., Littlewood, J. E., *Some more theorems concerning Fourier series and Fourier power series*, Duke Math. J. **2** (1936), no. 2, 354–381.
148. Hardy, G. H., Littlewood, J. E., Pólya, G., *Inequalities*, 2nd edition, Cambridge University Press, Cambridge, 1952.
149. Helson, H., Szegő, G., *A problem in prediction theory*, Ann. Mat. Pura Appl. (4) **51** (1960), 107–138.
150. Heo, Y., Nazarov, F., Seeger, A., *Radial Fourier multipliers in high dimensions*, Acta Math. **206** (2011), no. 1, 55–92.
151. Hernández, E., Weiss, G., *A First Course on Wavelets (Studies in Advanced Mathematics)*, CRC Press, Boca Raton, FL, 1996.
152. Hewitt, E., Ross, K., *Abstract Harmonic Analysis, Vol I*, 2nd edition, Grundlehren der mathematischen Wissenschaften, 115, Springer-Verlag, Berlin, New York, 1979.
153. Hewitt, E., Stromberg, K., *Real and Abstract Analysis*, Graduate Texts in Math. 25, Springer-Verlag, New York, 1965.
154. Hirschman, I. I. Jr., *A convexity theorem for certain groups of transformations*, J. Analyse Math. **2** (1953), 209–218.
155. Hofmann, S., *Weak $(1,1)$ boundedness of singular integrals with nonsmooth kernel*, Proc. Amer. Math. Soc. **103** (1988), no. 1, 260–264.

156. Hollenbeck, B., Verbitsky, I. E., *Best constants for the Riesz projection*, J. Funct. Anal. **175** (2000), no. 2, 370–392.
157. Honzík, P., *An example of an unbounded maximal singular operator*, J. Geom. Anal. **20** (2010), 153–167.
158. Honzík, P., *On p dependent boundedness of singular integral operators*, Math. Z. **267** (2011), no. 3–4, 931–937.
159. Hörmander, L., *Estimates for translation invariant operators in L^p spaces*, Acta Math. **104** (1960), no. 1–2, 93–140.
160. Hörmander, L., *The Analysis of Linear Partial Differential Operators I*, 2nd edition, Springer-Verlag, Berlin, 1990.
161. Hrusčev, S. V., *A description of weights satisfying the A_∞ condition of Muckenhoupt*, Proc. Amer. Math. Soc. **90** (1984), no. 2, 253–257.
162. Hudson, S., *A covering lemma for maximal operators with unbounded kernels*, Michigan Math. J. **34** (1987), no. 1, 147–151.
163. Hunt, R., *An extension of the Marcinkiewicz interpolation theorem to Lorentz spaces*, Bull. Amer. Math. Soc. **70** (1964), 803–807.
164. Hunt, R., *On $L(p, q)$ spaces*, Enseignement Math. (2) **12** (1966), 249–276.
165. Hunt, R., *On the convergence of Fourier series*, Orthogonal Expansions and Their Continuous Analogues (Proc. Conf., Edwardsville, IL, 1967), pp. 235–255, Southern Illinois Univ. Press, Carbondale IL, 1968.
166. Hunt, R., Kurtz, D., Neugebauer, C. J., *A note on the equivalence of A_p and Sawyer's condition for equal weights*, Conference on Harmonic Analysis in Honor of Antoni Zygmund, Vol. I, II (Chicago, IL, 1981), pp. 156–158, Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983.
167. Hunt, R., Muckenhoupt, B., Wheeden, R., *Weighted norm inequalities for the conjugate function and the Hilbert transform*, Trans. Amer. Math. Soc. **176** (1973), 227–251.
168. Hytönen, T. P., *The sharp weighted bound for general Calderón-Zygmund operators*, Ann. of Math. (2nd Ser.) **175** (2012), no.3, 1473–1506.
169. Hytönen, T. P., Lacey, M. T., Pérez, C., *Sharp weighted bounds for the q -variation of singular integrals*, Bull. Lond. Math. Soc. **45** (2013), no. 3, 529–540.
170. Hytönen, T. P., Pérez, C., *Sharp weighted bounds involving A_∞* , Anal. PDE **6** (2013), no. 4, 777–818.
171. Igari, S., *Lectures on Fourier Series of Several Variables*, Univ. Wisconsin Lecture Notes, Madison, WI, 1968.
172. Ionescu, A. D., *An endpoint estimate for the Kunze-Stein phenomenon and related maximal operators*, Ann. Math. (2) **152** (2000), no. 1, 259–275.
173. Iosevich, A., Sawyer, E., *Maximal averages over surfaces*, Adv. Math. **132** (1997), no. 1, 46–119.
174. Ismagilov, R. S., *On the Pauli problem*, Funct. Anal. Appl. **30** (1996), no. 2, 138–140.
175. Iwaniec, T., Martin, G., *Riesz transforms and related singular integrals*, J. Reine Angew. Math. **473** (1996), 25–57.
176. Jessen, B., Marcinkiewicz, J., Zygmund, A., *Note on the differentiability of multiple integrals*, Fund. Math. **25** (1935), 217–234.
177. Jodeit, M. Jr., *Restrictions and extensions of Fourier multipliers*, Studia Math. **34** (1970), 215–226.
178. Jodeit, M., *A note on Fourier multipliers*, Proc. Amer. Math. Soc. **27** (1971), 423–424.
179. Jones, P., *Factorization of A_p weights*, Ann. of Math. (2nd Ser.) **111** (1980), no. 3, 511–530.
180. Jordan, C., *Sur la série de Fourier*, Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences **92** (1881), 228–230.
181. Journé, J.-L., *Calderón-Zygmund operators on product spaces*, Rev. Mat. Iberoamericana **1** (1985), no. 3, 55–91.
182. Kahane, J.-P., *The heritage of Fourier*, Perspectives in Analysis, 83–95, Math. Phys. Stud., **27**, Springer, Berlin, 2005.
183. Kahane, J.-P., Katznelson, Y., *Sur les ensembles de divergence des séries trigonométriques*, Studia Math. **26** (1966), 305–306.

184. Kaiser, G., *A Friendly Guide to Wavelets*, Birkhäuser, Boston, MA, 1994.
185. Kalton, N. J., *Convexity, type, and the three space problem*, *Studia Math.* **69** (1980/81), no. 3, 247–287.
186. Kalton, N. J., *Plurisubharmonic functions on quasi-Banach spaces*, *Studia Math.* **84** (1986), no. 3, 297–323.
187. Kalton, N. J., *Analytic functions in non-locally convex spaces and applications*, *Studia Math.* **83** (1986), no. 3, 275–303.
188. Kalton, N. J., Peck, N. T., Roberts, J. W., *An F -space Sampler*, London Mathematical Society Lecture Notes Series, 89, Cambridge University Press, Cambridge, 1984.
189. Katznelson, Y., *Sur les ensembles de divergence des séries trigonométriques*, *Studia Math.* **26** (1966), 301–304.
190. Katznelson, Y., *An Introduction to Harmonic Analysis*, 2nd corrected edition, Dover Publications, Inc., New York, 1976.
191. Kendall, D. G., *On the number of lattice points inside a random oval*, *Quart. J. Math., Oxford Ser.* **19** (1948), 1–26.
192. Kenig, C., Tomas, P. A., *Maximal operators defined by Fourier multipliers*, *Studia Math.* **68** (1980), no. 1, 79–83.
193. Khintchine, A., *Über dyadische Brüche*, *Math. Z.* **18** (1923), no. 1, 109–116.
194. Kislyakov, S., Kruglyak, N., *Extremal Problems in Interpolation Theory, Whitney–Besicovitch Coverings, and Singular Integrals*, Birkhäuser/Springer Basel AG, Basel, 2013.
195. Kolmogorov, A. N., *Une série de Fourier–Lebesgue divergente presque partout*, *Fund. Math.* **4** (1923), 324–328.
196. Kolmogorov, A. N., *Une contribution à l'étude de la convergence des séries de Fourier*, *Fund. Math.* **5** (1924), 96–97.
197. Kolmogorov, A. N., *Sur les fonctions harmoniques conjuguées et les séries de Fourier*, *Fund. Math.* **7** (1925), 23–28.
198. Kolmogorov, A. N., *Une série de Fourier–Lebesgue divergente partout*, *C. R. Acad. Sci. Paris* **183** (1926), 1327–1328.
199. Kolmogorov, A. N., *Zur Normierbarkeit eines topologischen Raumes*, *Studia Math.* **5** (1934), 29–33.
200. Konyagin, S. V., *On the divergence everywhere of trigonometric Fourier series*, *Sb. Math.* **191** (2000), no. 1–2, 97–120.
201. Korenovskyy, A. A., Lerner, A. K., Stokolos, A. M., *A note on the Gurov–Reshetnyak condition*, *Math. Res. Lett.* **9** (2002), no. 5–6, 579–583.
202. Körner, T. W., *Fourier Analysis*, Cambridge University Press, Cambridge, 1988.
203. Krantz, S. G., *A panorama of Harmonic Analysis*, Carus Mathematical Monographs, 27, Mathematical Association of America, Washington, DC, 1999.
204. Krantz, S. G., Parks, H. R., *The Geometry of Domains in Space*, Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser Boston, Inc., Boston, MA, 1999.
205. Krantz, S. G., Parks, H. R., *A Primer of Real Analytic Functions*, 2nd edition, Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser Boston, Inc., Boston, MA, 2002.
206. Krivine, J. L., *Théorèmes de factorisation dans les espaces réticulés*, Séminaire Maurey–Schwartz 1973–74: Espaces L^p , applications radonifiantes et géométrie des espaces de Banach, Exp. Nos. 22 et 23, Centre de Math., École Polytech., Paris, 1974.
207. Kruglyak, N., *An elementary proof of the real version of the Riesz–Thorin theorem*, *Interpolation Theory and Applications*, pp. 179–182, *Contemp. Math.* **445**, Amer. Math. Soc., Providence, RI, 2007.
208. Kurtz, D., *Operator estimates using the sharp function*, *Pacific J. Math.* **139** (1989), no. 2, 267–277.
209. Kurtz, D., Wheeden, R., *Results on weighted norm inequalities for multipliers*, *Trans. Amer. Math. Soc.* **255** (1979), 343–362.
210. Lacey, M. T., Petermichl, S., Reguera, M. C., *Sharp A_2 inequality for Haar shift operators*, *Math. Ann.* **348** (2010), no. 1, 127–141.
211. Laeng, E., *On the L^p norms of the Hilbert transform of a characteristic function*, *J. Funct. Anal.* **262** (2012), no. 10, 4534–4539.

212. Landau, E., *Zur analytischen Zahlentheorie der definiten quadratischen Formen (Über die Gitterpunkte in einem mehrdimensionalen Ellipsoid)*, Berl. Sitzungsber. **31** (1915), 458–476; reprinted in E. Landau “Collected Works,” Vol. 6, pp. 200–218, Thales-Verlag, Essen, 1986.
213. Lang, S., *Real Analysis*, 2nd edition, Addison-Wesley Publishing Company, Advanced Book Program, Reading, MA, 1983.
214. Lebesgue, H., *Intégrale, longueur, aire*, Annali Mat. Pura Appl. **7** (1902), 231–359.
215. Lebesgue, H., *Oeuvres Scientifiques (en cinq volumes)*, Vol. I, (French) Sous la rédaction de F. Châtelet et G. Choquet, Institut de Mathématiques de l’ Université de Genève, Geneva, 1972.
216. Lemarié, P., Meyer, Y., *Ondelettes et bases hilbertiennes*, Rev. Mat. Iberoamericana **2** (1986), no. 1–2, 1–18.
217. Lerner, A. K., *On pointwise estimates for maximal and singular integral operators*, Studia Math. **138** (2000), no. 3, 285–291.
218. Lerner, A. K., *An elementary approach to several results on the Hardy–Littlewood maximal operator*, Proc. Amer. Math. Soc., **136** (2008), no. 8, 2829–2833.
219. Lerner, A. K., *A pointwise estimate for the local sharp maximal function with applications to singular integrals*, Bull. London Math. Soc. **42** (2010), no. 5, 843–856.
220. Lerner, A. K., *On an estimate of Calderón–Zygmund operators by dyadic positive operators*, J. Anal. Math. **121** (2013), 141–161.
221. Lerner, A. K., *A simple proof of the A_2 conjecture*, Int. Math. Res. Not. IMRN **14** (2013), 3159–3170.
222. Lerner, A. K., *Mixed A_p – A_r inequalities for classical singular integrals and Littlewood–Paley operators*, J. Geom. Anal. **23** (2013), no. 3, 1343–1354.
223. Lerner, A. K., Ombrosi, S., Pérez, C., *A_1 bounds for Calderón–Zygmund operators related to a problem of Muckenhoupt and Wheeden*, Math. Res. Lett. **16** (2009), no. 1, 149–156.
224. Liang, Y. Y., Liu, L. G., Yang, D. C., *An Off-diagonal Marcinkiewicz interpolation theorem on Lorentz spaces*, Acta Math. Sin. (Engl. Ser.) **27** (2011), no. 8, 1477–1488.
225. Lions, J.-L., Peetre, J., *Sur une classe d’espaces d’ interpolation*, Inst. Hautes Études Sci. Publ. Math. No. **19** (1964), 5–68.
226. Littlewood, J. E., *On a certain bilinear form*, Quart. J. Math. Oxford Ser. **1** (1930), 164–174.
227. Littlewood, J. E., Paley, R. E. A. C., *Theorems on Fourier series and power series*, J. London Math. Soc. **6** (1931), no. 3, 230–233.
228. Littlewood, J. E., Paley, R. E. A. C., *Theorems on Fourier series and power series (II)*, Proc. London Math. Soc. **42** (1936), no. 1, 52–89.
229. Littlewood, J. E., Paley, R. E. A. C., *Theorems on Fourier series and power series (III)*, Proc. London Math. Soc. **43** (1937), no. 2, 105–126.
230. Loomis, L. H., *A note on the Hilbert transform*, Bull. Amer. Math. Soc. **52** (1946), 1082–1086.
231. Loomis, L. H., Whitney, H., *An inequality related to the isoperimetric inequality*, Bull. Amer. Math. Soc. **55** (1949), 961–962.
232. Lorentz, G. G., *Some new function spaces*, Ann. of Math. (2nd Ser.) **51** (1950), no. 1, 37–55.
233. Lorentz, G. G., *On the theory of spaces Λ* , Pacific. J. Math. **1** (1951), 411–429.
234. Lu, S., Ding, Y., Yan, D., *Singular Integrals and Related Topics*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
235. Luzin, N., *Sur la convergence des séries trigonométriques de Fourier*, C. R. Acad. Sci. Paris **156** (1913), 1655–1658.
236. Lyapunov, A., *Sur les fonctions-vecteurs complètement additives* [Russian], Izv. Akad. Nauk SSSR Ser. Mat. **4** (1940), 465–478.
237. Magyar, A., Stein, E. M., Wainger, S., *Discrete analogues in harmonic analysis: spherical averages*, Ann. of Math. (2nd Ser.) **155** (2002), no. 1, 189–208.
238. Mallat, S., *Multiresolution approximations and wavelet orthonormal bases of $L^2(\mathbb{R})$* , Trans. Amer. Math. Soc. **315** (1989), no. 1, 69–87.
239. Mallat, S., *A Wavelet Tour of Signal Processing*, Academic Press, Inc., San Diego, CA, 1998.
240. Marcinkiewicz, J., *Sur l’interpolation d’operations*, C. R. Acad. Sci. Paris **208** (1939), 1272–1273.

241. Marcinkiewicz, J., *Sur les multiplicateurs des séries de Fourier*, *Studia Math.* **8** (1939), 78–91.
242. Marcinkiewicz, J., Zygmund, A., *Quelques inégalités pour les opérations linéaires*, *Fund. Math.* **32** (1939), 112–121.
243. Marcinkiewicz, J., Zygmund, A., *On the summability of double Fourier series*, *Fund. Math.* **32** (1939), 122–132.
244. Mateu, J., Orobitg, J., Verdera, J., *Estimates for the maximal singular integral in terms of the singular integral: the case of even kernels*, *Ann. of Math. (2nd Ser.)* **174** (2011), no. 3, 1429–1483.
245. Mateu, J., Verdera, J., *L^p and weak L^1 estimates for the maximal Riesz transform and the maximal Beurling transform*, *Math. Res. Lett.* **13** (2006), no. 5–6, 957–966.
246. Mattila, P., *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, *Cambridge Studies in Advanced Mathematics*, 44, Cambridge University Press, Cambridge, 1995.
247. Maurey, B., *Théorèmes de factorization pour les opérateurs linéaires à valeurs dans les espace L^p* , *Astérisque*, No. 11, Société Mathématique de France, Paris, 1974.
248. Melas, A., *The best constant for the centered Hardy–Littlewood maximal inequality*, *Ann. of Math. (2nd Ser.)* **157** (2003), no. 2, 647–688.
249. Meyer, Y., *Principe d’incertitude, bases hilbertiennes et algèbres d’opérateurs*, *Séminaire Bourbaki*, Vol. 1985/86, *Astérisque* No. 145–146 (1987) **4**, 209–223.
250. Meyer, Y., *Ondelettes et Opérateurs, I, Ondelletes*, Hermann, Paris, 1990.
251. Meyer, Y., *Ondelettes et Opérateurs, II, Opérateurs de Calderón-Zygmund*, Hermann, Paris, 1990.
252. Meyer, Y., *Wavelets, Vibrations, and Scalings*, CRM Monograph Series, 9, American Mathematical Society, Providence, RI, 1998.
253. Meyer, Y., Coifman, R. R., *Ondelettes et Opérateurs, III, Opérateurs multilinéaires*, Hermann, Paris, 1991.
254. Mihlin, S. G., *On the multipliers of Fourier integrals* [Russian], *Dokl. Akad. Nauk. SSSR (N.S.)* **109** (1956), 701–703.
255. Miyachi, A., *On some singular Fourier multipliers*, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **28** (1981), no. 2, 267–315.
256. Mockenhaupt, G., Seeger, A., Sogge, C., *Wave front sets, local smoothing and Bourgain’s circular maximal theorem*, *Ann. of Math. (2nd Ser.)* **136** (1992), no. 1, 207–218.
257. Moon, K. H., *On restricted weak type $(1,1)$* , *Proc. Amer. Math. Soc.* **42** (1974), 148–152.
258. Morse, A. P., *Perfect blankets*, *Trans. Amer. Math. Soc.* **69** (1947), 418–442.
259. Muckenhoupt, B., *On certain singular integrals*, *Pacific J. Math.* **10** (1960), 239–261.
260. Muckenhoupt, B., *Weighted norm inequalities for the Hardy maximal function*, *Trans. Amer. Math. Soc.* **165** (1972), 207–226.
261. Muckenhoupt, B., *The equivalence of two conditions for weight functions*, *Studia Math.* **49** (1973/1974), 101–106.
262. Muckenhoupt, B., Wheeden, R. L., *Two weight function norm inequalities for the Hardy–Littlewood maximal function and the Hilbert transform*, *Studia Math.* **55** (1976), no. 3, 279–294.
263. Müller, D., *A geometric bound for maximal functions associated to convex bodies*, *Pacific J. Math.* **142** (1990), no. 2, 297–312.
264. Nagel, A., Stein, E. M., Wainger, S., *Differentiation in lacunary directions*, *Proc. Nat. Acad. Sci. U.S.A.* **75** (1978), no. 3, 1060–1062.
265. Nazarov, F., Reznikov, A., Vasyunin, V., Volberg, A., *A_1 conjecture: weak norm estimates of weighted singular integrals and Bellman functions*, <http://sashavolberg.wordpress.com>
266. Nazarov, F., Treil, S., Volberg, A., *The Bellman functions and two-weight inequalities for Haar multipliers*, *J. Amer. Math. Soc.* **12** (1999), no. 4, 909–928.
267. Nazarov, F., Treil, S., Volberg, A., *Bellman function in stochastic control and harmonic analysis*, *Systems, Approximation, Singular Integral Operators, and Related Topics* (Bordeaux, 2000), pp. 393–423, *Oper. Theory Adv. Appl.* 129, Birkhäuser, Basel, 2001.

268. Nazarov, F., Treil, S., Volberg, A., *Two weight inequalities for individual Haar multipliers and other well localized operators*, Math. Res. Lett. **15** (2008), no. 3, 583–597.
269. Nguyen, H. V., *A maximal Fourier integral operator and an application*, J. of Pseudodifferential Operators **4** (2013), 443–456.
270. Nyquist, H., *Certain topics in telegraph transmission theory*, Trans. AIEE **47** (Apr. 1928), 617–64.
271. Olver, F. W. J., *Asymptotics and Special Functions*, Academic Press, New York, London, 1974.
272. Orobitg, J., Pérez, C., *A_p weights for nondoubling measures in R^n and applications*, Trans. Amer. Math. Soc. **354** (2002), no. 5, 2013–2033 (electronic).
273. Paley, R. E. A. C., *A remarkable series of orthogonal functions (II)*, Proc. London Math. Soc. **34** (1932), 241–264.
274. Paley, R. E. A. C., Zygmund, A., *On some series of functions*, Proc. Cambridge Phil. Soc. **26** (1930), 337–357.
275. Peetre, J., *Nouvelles propriétés d'espaces d'interpolation*, C. R. Acad. Sci. Paris **256** (1963), 1424–1426.
276. Pereyra, M. C., *Lecture Notes on Dyadic Harmonic Analysis*, Second Summer School in Analysis and Mathematical Physics (Cuernavaca, 2000), pp. 1–60, Contemp. Math., 289, American Mathematical Society, Providence, RI, 2001.
277. Pérez, C., *Weighted norm inequalities for singular integral operators*, J. London Math. Soc. (2) **49** (1994), no. 2, 296–308.
278. Pérez, C., *Some topics from Calderón–Zygmund theory related to Poincaré–Sobolev inequalities, fractional integrals and singular integral operators*, Function Spaces Lectures, Spring School in Analysis, pp. 31–94, Jaroslav Lukeš and Luboš Pick (eds.), Paseky, 1999.
279. Petermichl, S., *The sharp bound for the Hilbert transform on weighted Lebesgue spaces in terms of the classical A_p characteristic*, Amer. J. Math. **129** (2007), no. 5, 1355–1375.
280. Petermichl, S., *The sharp weighted bound for the Riesz transforms*, Proc. Amer. Math. Soc. **136** (2008), no. 4, 1237–1249.
281. Petermichl, S., Volberg, A., *Heating of the Ahlfors–Beurling operator: weakly quasiregular maps on the plane are quasiregular*, Duke Math. J. **112** (2002), no. 2, 281–305.
282. Pichorides, S., *On the best values of the constants in the theorems of M. Riesz, Zygmund and Kolmogorov*, Studia Math. **44** (1972), 165–179.
283. Pinsky, M., *Introduction to Fourier Analysis and Wavelets*, Graduate Studies in Mathematics 102, American Mathematical Society, Providence, RI, 2009.
284. Pinsky, M., Stanton, N. K., Trapa, P. E., *Fourier series of radial functions in several variables*, J. Funct. Anal. **116** (1993), no. 1, 111–132.
285. Plancherel, M., Pólya, G., *Fonctions entières et intégrales de fourier multiples*, Comment. Math. Helv. **10** (1937), no. 1, 110–163.
286. Reed, M., Simon, B., *Methods of Modern Mathematical Physics, II, Fourier analysis, self-adjointness*, Academic Press, New York, London, 1975.
287. Reguera, M. C., *On Muckenhoupt–Wheeden conjecture*, Adv. Math. **227** (2011), no. 4, 1436–1450.
288. Reguera, M., Thiele, C., *The Hilbert transform does not map $L^1(Mw)$ to $L^{1,\infty}(w)$* , Math. Res. Lett. **19** (2012), no. 1, 1–7.
289. Ricci, F., Weiss, G., *A characterization of $H^1(\Sigma_{n-1})$* , Harmonic Analysis in Euclidean Spaces Proc. Sympos. Pure Math. (Williamstown, MA, 1978), Part 1, pp. 289–294, Amer. Math. Soc., Providence, RI, 1979.
290. Riesz, F., *Untersuchungen über Systeme integrierbarer Funktionen*, Math. Ann. **69** (1910), no. 4, 449–497.
291. Riesz, F., *Sur un théorème du maximum de Mm. Hardy et Littlewood*, J. London Math. Soc. **7** (1932), 10–13.
292. Riesz, M., *Les fonctions conjuguées et les séries de Fourier*, C. R. Acad. Sci. Paris **178** (1924), 1464–1467.
293. Riesz, M., *Sur les maxima des formes bilinéaires et sur les fonctionnelles linéaires*, Acta Math. **49** (1927), no. 3–4, 465–497.

294. Riesz, M., *Sur les fonctions conjuguées*, Math. Z. **27** (1928), no. 1, 218–244.
295. Riesz, M., *L'intégrale de Riemann–Liouville et le problème de Cauchy*, Acta Math. **81** (1949), no. 1, 1–223.
296. Riviere, N., *Singular integrals and multiplier operators*, Ark. Mat. **9** (1971), 243–278.
297. Rolewicz, S., *Metric Linear Spaces*, 2nd edition, Mathematics and its Applications (East European Series), 20. D. Reidel Publishing Co., Dordrecht; PWN–Polish Scientific Publishers, Warsaw, 1985.
298. Rosenblum, M., *Summability of Fourier series in $L^p(d\mu)$* , Trans. Amer. Math. Soc. **105** (1962), 32–42.
299. Rubio de Francia, J.-L., *Weighted norm inequalities and vector valued inequalities*, Harmonic Analysis (Minneapolis, MN, 1981), pp. 86–101, Lect. Notes in Math. 908, Springer, Berlin, New York, 1982.
300. Rubio de Francia, J.-L., *Factorization theory and A_p weights*, Amer. J. Math. **106** (1984), no. 3, 533–547.
301. Rubio de Francia, J.-L., *A Littlewood–Paley inequality for arbitrary intervals*, Rev. Mat. Iberoamericana **1** (1985), no. 2, 1–14.
302. Rubio de Francia, J.-L., *Maximal functions and Fourier transforms*, Duke Math. J. **53** (1986), no. 2, 395–404.
303. Rubio de Francia, J.-L., Ruiz, F. J., Torrea, J. L., *Calderón–Zygmund theory for operator-valued kernels*, Adv. in Math. **62** (1986), no. 1, 7–48.
304. Rubio de Francia, J.-L., Torrea, J. L., *Vector extensions of operators in L^p spaces*, Pacific J. Math. **105** (1983), no. 1, 227–235.
305. Rudin, W., *Trigonometric series with gaps*, J. Math. Mech. **9** (1960), 203–227.
306. Rudin, W., *Functional Analysis*, Tata McGraw-Hill Publishing Company, New Delhi, 1973.
307. Rudin, W., *Real and Complex Analysis*, 2nd edition, Tata McGraw-Hill Publishing Company, New Delhi, 1974.
308. Ryabogin, D., Rubin, B., *Singular integrals generated by zonal measures*, Proc. Amer. Math. Soc. **130** (2002), no. 3, 745–751.
309. Sadosky, C., *Interpolation of Operators and Singular Integrals*, Marcel Dekker Inc., New York, 1979.
310. Saks, S., *Theory of the Integral*, Hafner Publ. Co, New York, 1938.
311. Sato, S., *Note on a Littlewood–Paley operator in higher dimensions*, J. London Math. Soc. (2) **42** (1990), no. 3, 527–534.
312. Sawyer, E., *A characterization of a two-weight norm inequality for maximal operators*, Studia Math. **75** (1982), no. 1, 1–11.
313. Schlag, W., *A geometric proof of the circular maximal theorem*, Duke Math. J. **93** (1998), no. 3, 505–533.
314. Schwartz, L., *Théorie de Distributions. Tome I*, Hermann & Cie, Paris, 1950.
315. Schwartz, L., *Théorie de Distributions. Tome II*, Hermann & Cie, Paris, 1951.
316. Seeger, A., *Some inequalities for singular convolution operators in L^p -spaces*, Trans. Amer. Math. Soc. **308** (1988), no. 1, 259–272.
317. Seeger, A., *Singular integral operators with rough convolution kernels*, J. Amer. Math. Soc. **9** (1996), no. 1, 95–105.
318. Seeger, A., Tao, T., *Sharp Lorentz estimates for rough operators*, Math. Ann. **320** (2001), no. 2, 381–415.
319. Shannon, C. E., *Communication in the presence of noise*, Proc. I.R.E. **37** (1949), 10–21.
320. Shapiro, V. L., *Fourier series in several variables*, Bull. Amer. Math. Soc. **70** (1964), 48–93.
321. Shapiro, V. L., *Fourier series in several variables with applications to partial differential equations*, Chapman & Hall/CRC Applied Mathematics and Nonlinear Science Series, CRC Press, Boca Raton, FL, 2011.
322. Siegel, A., *Historical review of the isoperimetric theorem in 2-D, and its place in elementary plane geometry*, <http://www.cs.nyu.edu/faculty/siegel/SCIAM.pdf>
323. Sjögren, P., *A remark on the maximal function for measures in \mathbb{R}^n* , Amer. J. Math. **105** (1983), no. 5, 1231–1233

324. Sjögren, P., Soria, F., *Rough maximal operators and rough singular integral operators applied to integrable radial functions*, Rev. Math. Iberoamericana **13** (1997), no. 1, 1–18.
325. Sjölin, P., *Convergence almost everywhere of certain singular integrals and multiple Fourier series*, Ark. Math. **9** (1971), 65–90.
326. Sjölin, P., *A note on Littlewood–Paley decompositions with arbitrary intervals*, J. Approx. Theory **48** (1986), no. 3, 328–334.
327. Sjölin, P., Soria, F., *Remarks on a theorem by N. Yu. Antonov*, Studia Math. **158** (2003), no. 1, 79–97.
328. Sogge, C., *Fourier Integrals in Classical Analysis*, Cambridge Tracts in Mathematics, 105, Cambridge University Press, Cambridge, 1993.
329. Soria, F., *A note on a Littlewood–Paley inequality for arbitrary intervals in \mathbb{R}^2* , J. London Math. Soc. (2) **36** (1987), no. 1, 137–142.
330. Soria, F., *On an extrapolation theorem of Carleson–Sjölin with applications to a.e. convergence of Fourier series*, Studia Math. **94** (1989), no. 3, 235–244.
331. Stein, E. M., *Interpolation of linear operators*, Trans. Amer. Math. Soc. **83** (1956), 482–492.
332. Stein, E. M., *Note on singular integrals*, Proc. Amer. Math. Soc. **8** (1957), 250–254.
333. Stein, E. M., *Localization and summability of multiple Fourier series*, Acta Math. **100** (1958), no. 1–2, 93–147.
334. Stein, E. M., *On the functions of Littlewood–Paley, Lusin, and Marcinkiewicz*, Trans. Amer. Math. Soc. **88** (1958), 430–466.
335. Stein, E. M., *On limits of sequences of operators*, Ann. of Math. (2nd Ser.) **74** (1961), no. 1, 140–170.
336. Stein, E. M., *Note on the class $L \log L$* , Studia Math. **32** (1969), 305–310.
337. Stein, E. M., *Topics in Harmonic Analysis Related to the Littlewood–Paley Theory*, Annals of Mathematics Studies, No. 63, Princeton University Press, Princeton, NJ, 1970.
338. Stein, E. M., *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, NJ, 1970.
339. Stein, E. M., *Maximal functions I: Spherical means*, Proc. Nat. Acad. Sci. U.S.A. **73** (1976), no. 7, 2174–2175.
340. Stein, E. M., *The development of square functions in the work of A. Zygmund*, Bull. Amer. Math. Soc. (N.S.) **7** (1982), no. 2, 359–376.
341. Stein, E. M., *Some results in harmonic analysis in \mathbb{R}^n , for $n \rightarrow \infty$* , Bull. Amer. Math. Soc. (N.S.) **9** (1983), no. 1, 71–73.
342. Stein, E. M., *Boundary behavior of harmonic functions on symmetric spaces: Maximal estimates for Poisson integrals*, Invent. Math. **74** (1983), no. 1, 63–83.
343. Stein, E. M., *Oscillatory integrals in Fourier analysis*, Beijing Lectures in Harmonic Analysis (Beijing, 1984), pp. 307–355, Annals of Math. Studies 112, Princeton Univ. Press, Princeton, NJ, 1986.
344. Stein, E. M., *Harmonic Analysis, Real Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series, 43, Monographs in Harmonic Analysis, III, Princeton University Press, Princeton, NJ, 1993.
345. Stein, E. M., Strömberg, J.-O., *Behavior of maximal functions in \mathbb{R}^n for large n* , Ark. Math. **21** (1983), no. 2, 259–269.
346. Stein, E. M., Weiss, G., *Interpolation of operators with change of measures*, Trans. Amer. Math. Soc. **87** (1958), 159–172.
347. Stein, E. M., Weiss, G., *An extension of theorem of Marcinkiewicz and some of its applications*, J. Math. Mech. **8** (1959), 263–284.
348. Stein, E. M., Weiss, G., *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Mathematical Series, No. 32, Princeton University Press, Princeton, NJ, 1971.
349. Stein, E. M., Weiss, N. J., *On the convergence of Poisson integrals*, Trans. Amer. Math. Soc. **140** (1969), 34–54.
350. Stein, P., *On a theorem of M. Riesz*, J. London Math. Soc. **8** (1933), no. 4, 242–247.
351. Stepanov, V. D., *Convolution integral operators [Russian]*, Dokl. Akad. Nauk SSSR **243** (1978), no. 1, 45–48.

352. Strömberg, J.-O., *A modified Franklin system and higher-order spline systems on \mathbb{R}^n as unconditional bases for Hardy spaces*, Conference on Harmonic Analysis in Honor of Antoni Zygmund, Vol. I, II (Chicago, IL, 1981), pp. 475–494, Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983.
353. Szarek, S. J., *On the best constant in the Khinchin inequality*, *Studia Math.* **58** (1976), no. 2, 197–208.
354. Tamarkin, J. D., Zygmund, A., *Proof of a theorem of Thorin*, *Bull. Amer. Math. Soc.* **50** (1944), 279–282.
355. Tao, T., *The weak type (1,1) of $L \log L$ homogeneous convolution operators*, *Indiana Univ. Math. J.* **48** (1999), no. 4, 1547–1584.
356. Tao, T., *A converse extrapolation theorem for translation-invariant operators*, *J. Funct. Anal.* **180** (2001), no. 1, 1–10.
357. Tao, T., Wright, J., *Endpoint multiplier theorems of Marcinkiewicz type*, *Rev. Mat. Iberoamericana* **17** (2001), no. 3, 521–558.
358. Tauber, A., *Ein Satz aus der Theorie der unendlichen Reihen* *Monatsh. Math. Phys.* **8** (1897), no. 1, 273–277.
359. Tevzadze, N. R., *The convergence of the double Fourier series of quadratic at a square summable function*, *Sakharth. SSR Mecn. Akad. Moambe* **58** (1970), 277–279.
360. Thorin, G. O., *An extension of a convexity theorem due to M. Riesz*, *Fys. Sällsk. Förh.* **8** (1938), No. 14.
361. Thorin, G. O., *Convexity theorems generalizing those of M. Riesz and Hadamard with some applications*, *Comm. Sem. Math. Univ. Lund [Medd. Lunds Univ. Mat. Sem.]* **9** (1948), 1–58.
362. Titchmarsh, E. C., *The Theory of the Riemann Zeta Function*, Oxford, at the Clarendon Press, 1951.
363. Torchinsky, A., *Real-Variable Methods in Harmonic Analysis*, Pure and Applied Mathematics, 123, Academic Press, Inc., Orlando, FL, 1986.
364. van der Corput, J. G., *Zahlentheoretische Abschätzungen*, *Math. Ann.* **84** (1921), no. 1–2, 53–79.
365. Vargas, A., *On the maximal function for rotation invariant measures in \mathbb{R}^n* , *Studia Math.* **110** (1994), no. 1, 9–17.
366. Verbitsky, I. E., *Estimate of the norm of a function in a Hardy space in terms of the norms of its real and imaginary part*, *Amer. Math. Soc. Transl.* **124** (1984), 11–15.
367. Verbitsky, I. E., *Weighted norm inequalities for maximal operators and Pisier's theorem on factorization through L^{∞}* , *Integral Equations Operator Theory* **15** (1992), no. 1, 124–153.
368. Vitali, G., *Sui gruppi di punti e sulle funzioni di variabili reali*, *Atti Accad. Sci. Torino* **43** (1908), 229–246.
369. Von Neuman, J., *Zur Theorie des Gesellschaftsspiele*, *Math. Ann.* **100** (1928), 295–320.
370. Wainger, S., *Special trigonometric series in k dimensions*, *Mem. Amer. Math. Soc.* **59**, 1965.
371. Watson, G. N., *A Treatise on the Theory of Bessel Functions*, Cambridge University Press, Cambridge, England, The Macmillan Company, New York, 1944.
372. Weiss, M., Zygmund, A., *An example in the theory of singular integrals*, *Studia Math.* **26** (1965), 101–111.
373. Whitney, H., *Analytic extensions of differentiable functions defined in closed sets*, *Trans. Amer. Math. Soc.* **36** (1934), no. 1, 63–89.
374. Wickerhauser, M. V., *Adapted Wavelet Analysis from Theory to Software*, A. K. Peters, Ltd., Wellesley, MA, 1994.
375. Wiener, N., *The ergodic theorem*, *Duke Math. J.* **5** (1939), no. 1, 1–18.
376. Wilbraham, H., *On a certain periodic function*, *The Cambridge and Dublin Mathematical Journal*, **3** (1848), 198–201.
377. Wilson, J. M., *Weighted norm inequalities for the continuous square function*, *Trans. Amer. Math. Soc.* **314** (1989), no. 2, 661–692.
378. Wilson, M., *Weighted Littlewood–Paley Theory and Exponential-Square Integrability*, *Lecture Notes in Mathematics*, 1924 Springer, Berlin, 2008.
379. Wojtaszczyk, P., *A Mathematical Introduction to Wavelets*, London Mathematical Society Student Texts 37, Cambridge University Press, Cambridge, 1997.

380. Yano, S., *Notes on Fourier analysis, XXIX, An extrapolation theorem*, J. Math. Soc. Japan **3** (1951), 296–305.
381. Yanushauskas, A. I., *Multiple trigonometric series* [Russian], "Nauka" Sibirsk. Otdel., Novosibirsk, 1986.
382. Yosida, K., *Functional Analysis*, Reprint of the 6th edition (1980), Classics in Mathematics Springer-Verlag, Berlin, 1995.
383. Zhizhiashvili, L. V., *Some problems in the theory of simple and multiple trigonometric and orthogonal series*, Russian Math. Surveys, **28** (1973), 65–127.
384. Zhizhiashvili, L. V., *Trigonometric Fourier Series and Their Conjugates*, Mathematics and Its Applications, 372, Kluwer Academic Publishers Group, Dordrecht, 1996.
385. Ziemer, W. P., *Weakly Differentiable Functions, Sobolev Spaces and Functions of Bounded Variation*, Graduate Texts in Mathematics, 120, Springer-Verlag, New York, 1989.
386. Zo, F., *A note on approximation of the identity*, Studia Math. **55** (1976), no. 2, 111–122.
387. Zygmund, A., *On a theorem of Marcinkiewicz concerning interpolation of operators*, J. Math. Pures Appl. (9) **35** (1956), 223–248.
388. Zygmund, A., *Trigonometric Series*, Vol. I, 2nd edition, Cambridge University Press, New York, 1959.
389. Zygmund, A., *Trigonometric Series*, Vol. II, 2nd edition, Cambridge University Press, New York, 1959.
390. Zygmund, A., *Notes on the history of Fourier series*, Studies in harmonic analysis, (Proc. Conf. DePaul Univ., Chicago, IL, 1974) pp. 1–19, MAA Stud. Math., Vol. 13, Math. Assoc. Amer., Washington, DC, 1976.

Index

- \mathcal{C}^k functions on the torus, 175
- σ -finite, 2

- absolutely summable Fourier series, 200
- adjoint of an operator, 150
- admissible growth, 40
- almost everywhere convergence, 285
- almost orthogonality, 459, 471
- analytic family of operators, 40
- Aoki–Rolewicz theorem, 76
- A_1 condition, 502
- A_p condition, 503
 - necessity of, 543
- approximate identity, 26
- asymptotics
 - of Bessel function, 580
- asymptotics of gamma function, 567
- atom
 - in a measure space, 56

- bad function, 356
- Banach–Alaoglu theorem, 601
- Banach-valued extension of a linear operator, 391
- Banach-valued extension of an operator, 396
- Banach-valued measurable function, 392
- Banach-valued singular integral, 402
- band limited function, 490
- Bernoulli polynomials, 299
- Bernstein’s inequality, 133
- Bernstein’s theorem, 191
- Bessel function, 168, 573
 - asymptotics, 580
 - large arguments, 579
 - small arguments, 578
- beta function, 564
- beta integral identity, 145

- Boas and Bochner inequality, 391
- Bochner integral, 394
- Bochner–Riesz means, 244
- Bochner–Riesz operator, 244
- Borel measures, 2
- bounded variation, 199
- BV , functions of bounded variation, 199

- Calderón–Zygmund operator, 532
- Calderón–Zygmund decomposition, 356
- Calderón–Zygmund decomposition on L^q , 373
- Calderón–Zygmund operator
 - maximal, 533
- cancellation condition
 - for a kernel, 375
- Carleson operator, 286
- Cauchy sequence in measure, 9
- centered Hardy–Littlewood maximal function, 86
- centered maximal function with respect to
 - cubes, 99
- Cesàro means, 184, 216
- characteristic constant of a pair of weights, 524
- Characteristic constant of a weight
 - A_p , 503
- characteristic constant of a weight
 - A_1 , 503
- characterization
 - of A_1 weights, 519
 - of A_∞ weights, 527, 531
- Chebyshev’s inequality, 6
- circular Dirichlet kernel, 178
- circular means, 250
- circular partial sum, 184
- class (A_p, A_p) , 524
- class of A_1 weights, 502
- class of A_p weights, 503

- closed
 - under translations, 146
- closed graph theorem, 602
- commutes with translations, 146
- compactly supported distribution, 120
- complete orthonormal system, 185
- completeness
 - of Lorentz spaces, 54
- completeness of L^p , $p < 1$, 13
- conditional expectation, 464
- cone multiplier, 158
- conjugate function, 245
- conjugate harmonic, 318
- conjugate harmonic functions, 253
- conjugate Poisson kernel, 253, 318, 331
- continuously differentiable function
 - of order N , 104
- continuously differentiable functions
 - on the torus, 175
- convergence
 - in \mathcal{C}_0^∞ , 119
 - in \mathcal{S} , 106, 119
 - in L^p , 7
 - in \mathcal{C}^∞ , 119
 - in measure, 6
 - in weak L^p , 7
- convex sequence, 193
- convolution, 20, 125
- Cotlar's inequality, 364
- countably simple, 56
- countably simple function, 2
- countably subadditive operator, 47
- covering lemma, 88
- critical Bochner–Riesz index, 252
- critical point, 162

- Darboux's equation, 476
- de la Vallée Poussin kernel, 182
- decay of Fourier coefficients, 193
- decomposition of open sets, 609
- decreasing rearrangement, 48
 - of a simple function, 49
- degree of a trigonometric polynomial, 177
- deLeeuw's theorem, 157
- derivative
 - of a distribution, 123
 - of a function (partial), 104
- differentiation
 - of approximate identities, 96
 - theory, 93
- dilation
 - L^1 dilation, 90
 - of a function, 109
 - of a tempered distribution, 124
- Dini condition, 384
- Dini's theorem, 212
- Dirac mass, 121
- directional Hilbert transform, 339
- Dirichlet kernel, 178
- Dirichlet problem, 92
 - on the sphere, 144
- distribution, 120
 - homogeneous, 132, 136
 - of lattice points, 305
 - supported at a point, 134
 - tempered, 120
 - with compact support, 120, 129
- distribution function, 3
 - of a simple function, 4
- distributional derivative, 123
- divergence
 - of Bochner–Riesz means at the critical index, 261
 - of the Fourier series of a continuous function, 210
 - of the Fourier series of an L^1 function, 255
- doubling condition
 - on a measure, 98
- doubling measure, 98, 505
- doubly truncated kernel, 363
- doubly truncated singular integrals, 363
- duals of Lorentz spaces, 57
- duBois Reymond's theorem, 210
- duplication formula
 - for the gamma function, 570
- dyadic cube, 464
- dyadic decomposition, 428
- dyadic interval, 464
- dyadic martingale difference operator, 465
- dyadic martingale square function, 469
- dyadic maximal function, 103
- dyadic spherical maximal function, 461

- eigenvalues
 - of the Fourier transform, 116
- equidistributed, 48
- equidistributed sequence, 302
- essential infimum, 502
- Euler's constant, 569
- Euler's limit formula for the gamma function, 568
 - infinite product form, 569
- expectation
 - conditional, 464
- extrapolation, 47
- extrapolation theorem, 548, 553

- factorization of A_p weights, 546
- Fejér kernel, 27, 180

- Fejér means, 184
- Fejér's theorem, 205
- finite Borel measure, 2
- finitely simple function, 2
- Fourier coefficient, 175
 - of a measure, 175
- Fourier inversion, 112, 185
 - on L^1 , 117
 - on L^2 , 114
- Fourier multiplier, 155
- Fourier series, 176
- Fourier transform
 - of a radial function, 577
 - of a Schwartz function, 108
 - of surface measure, 577
 - on L^1 , 113
 - on L^2 , 113
 - on L^p , $1 < p < 2$, 114
 - properties of, 109
- Fréchet space, 106
- frame
 - tight, 496
- Fresnel integral, 145
- fundamental solution, 135
- fundamental theorem of algebra, 143

- g -function, 461, 477
- gamma function, 563
 - asymptotics, 567
 - duplication formula, 570
 - meromorphic extension, 566
- generalized Cauchy-Riemann equations, 332
- good function, 356, 372
- good lambda inequality, 533
 - consequences of, 539
- gradient, 104
- gradient condition
 - for a kernel, 359
- Green's identity, 135

- Hölder condition, 384
- Hölder's inequality, 3, 11
 - for weak spaces, 16
- Hörmander's condition, 359
- Hörmander–Mihlin multiplier theorem, 446
- Haar function, 465
- Haar measure, 18
- Haar wavelet, 483
- Hadamard's three lines lemma, 39
- Hahn–Banach theorem, 601
- Hardy's inequalities, 31
- Hardy–Littlewood maximal function
 - centered, 86
 - uncentered, 87
- harmonic distribution, 135
- harmonic function, 92
- Hausdorff–Young, 191
- Hausdorff–Young inequality, 114
- heat equation, 295
- heat kernel, 296
- Heisenberg group, 19
- Hilbert transform, 314
 - maximal, 322, 339
 - maximal directional, 339
 - truncated, 314
- Hirschman's lemma, 41
- homogeneous distribution, 132, 136
- homogeneous Lipschitz functions, 195
- homogeneous maximal singular integrals, 333
- homogeneous singular integrals, 333

- improved integrability
 - of A_p weights, 518
- inductive limit topology, 120
- infinitely differentiable function, 104
- inhomogeneous Lipschitz functions, 196
- inner product
 - complex, 150
 - real, 150
- inner regular measure, 98
- interpolation
 - Banach-valued Marcinkiewicz theorem, 400
 - Banach-valued Riesz–Thorin theorem, 399
 - for analytic families of operators, 41
 - Marcinkiewicz theorem, 33
 - off-diagonal Marcinkiewicz theorem, 61
 - Riesz–Thorin theorem, 37
 - Stein's theorem, 41
 - with change of measure, 78
- inverse Fourier transform, 112
- isoperimetric inequality, 17

- Jensen's inequality, 12, 525
- Jones's factorization theorem, 546

- Khintchine's inequalities, 586
 - for weak type spaces, 589
- Kolmogorov's inequality, 100
- Kolmogorov's theorem, 255
- Kronecker's lemma, 255

- lacunary Carleson–Hunt theorem, 454
- lacunary sequence, 227
- Laplace's equation, 328
- Laplacian, 135
- lattice points, 305
- Lebesgue constants, 182
- Lebesgue differentiation theorem, 95, 101

- left Haar measure, 18
- left maximal function, 102
- Leibniz's rule, 105, 131
- linear operator, 33
- Liouville's theorem, 143
- Lipschitz condition, 384
 - for a kernel, 359
- Lipschitz space
 - homogeneous, 195
 - inhomogeneous, 196
- Littlewood–Paley operator, 420
- Littlewood–Paley theorem, 421
- localization principle, 213
- locally compact topological group, 18
- locally finite measure, 98
- locally integrable functions, 11
- logconvexity of weak L^1 , 78
- Lorentz spaces, 52

- M. Riesz's theorem, 247
- majorant
 - radial decreasing, 92
- Marcinkiewicz and Zygmund theorem, 386
- Marcinkiewicz function, 415
- Marcinkiewicz interpolation theorem, 33
- Marcinkiewicz multiplier theorem
 - on \mathbf{R}^n , 441
 - on \mathbf{R} , 439
- maximal Calderón–Zygmund operator, 533
- maximal function
 - centered with respect to cubes, 99
 - dyadic, 103
 - dyadic spherical, 461
 - Hardy–Littlewood centered, 86
 - Hardy–Littlewood uncentered, 87
 - left, 102
 - right, 102
 - strong, 101
 - uncentered with respect to cubes, 99
 - with respect to a general measure, 98
- maximal Hilbert transform, 322
- maximal singular integral, 334
 - doubly truncated, 363
- maximal singular integrals with even kernels, 347
- maximal truncated singular integral, 363
- measurable set, 2
- method of rotations, 339
- metrizability
 - of Lorentz space $L^{p,q}$, 74
 - of weak L^p , 14
- Mihlin–Hörmander multiplier theorem, 446
- minimally supported frequency wavelet, 486
- minimax lemma, 556, 603

- Minkowski's inequality, 3, 12, 21
 - integral form, 13
- Minkowski's integral inequality, 13
- Muckenhoupt characteristic constant, 503
- Muckenhoupt characteristic constant of a weight, 503
- multi-index, 104
- multiplier, 155
 - on the torus, 272
- multiplier theorems, 437

- necessity of A_p condition, 543
- nonatomic measure space, 56
- nonsmooth Littlewood–Paley theorem, 427, 428
- normability
 - p -normability of weak L^p for $p < 1$, 16
 - of Lorentz space $L^{p,q}$, 74
 - of Lorentz spaces, 74
 - of weak L^p for $p > 1$, 14
- normability of weak L^1 (lack of), 15

- off-diagonal Marcinkiewicz interpolation theorem, 61
- open mapping theorem, 602
- operator
 - commuting with translations, 146
 - of strong type (p, q) , 33
 - of weak type (p, q) , 33
- orthonormal set, 185, 484
- orthonormal system
 - complete, 185
- oscillation of a function, 94
- oscillatory integral, 161
- overshoot, 221, 224

- Parseval's relation, 112, 186
- partial derivative, 104
- partition of unity, 611
- phase, 161
- Plancherel's identity, 112, 186
- pointwise convergence of Fourier series, 204
- Poisson kernel, 26, 92, 95, 183, 317
- Poisson kernel for the sphere, 144
- Poisson representation formula
 - of Bessel functions, 573
- Poisson summation formula, 187
- polarization, 186
- positive operator, 397
- power weights, 506
- principal value integral, 314
- principle of localization, 213

- quasi-linear operator, 33, 61
- quasi-subadditive operator, 45

- Rademacher functions, 585
- radial decreasing majorant, 92
- radial function, 90
- real analytic function, 262
- reflection
 - of a function, 109
 - of a tempered distribution, 124
- reflection formula
 - for the gamma function, 570
- regulated function, 275
- restricted weak type, 77
- restricted weak type $(1, 1)$, 81
- restricted weak type (p, q) with constant C , 61
- reverse Hölder condition
 - of order q , 523
- reverse Hölder property of A_p weights, 514
- $RH_q(\mu)$, 523
- Riemann zeta function, 301
- Riemann's principle of localization, 213
- Riemann–Lebesgue lemma, 114, 193, 215
- Riesz product, 232
- Riesz projection, 245
- Riesz transform, 325
- Riesz–Thorin interpolation theorem, 37
- right maximal function, 102
- right Haar measure, 18
- rotations method, 339
- rough singular integrals, 455
- Rubio de Francia's extrapolation theorem, 548

- sampling theorem, 491
- Schwartz function, 105
- Schwartz seminorm, 105
- self-adjoint operator, 151
- self-transpose operator, 151
- Sidon set, 235
- simple function, 2
- sine integral function, 218
- singular integrals with even kernels, 341
- size condition
 - for a kernel, 358, 374
- smooth bump, 120
- smooth function, 104, 119
- smooth function with compact support, 104, 119
- smoothly truncated maximal singular integral, 374
- smoothness condition
 - for a kernel, 359, 375
- space
 - L^∞ , 2
 - L^p , 2
 - $L^{p,\infty}$, 5
 - $L^{p,q}$, 52
 - $\mathcal{M}^{1,1}(\mathbf{R}^n)$, 153
 - $\mathcal{M}^{2,2}(\mathbf{R}^n)$, 154
 - $\mathcal{M}^{\infty,\infty}(\mathbf{R}^n)$, 154
 - $\mathcal{M}^{p,q}(\mathbf{R}^n)$, 151
 - $\mathcal{M}_p(\mathbf{R}^n)$, 155
 - \mathcal{E}^N , 104
 - \mathcal{E}^∞ , 104
 - \mathcal{E}_0 , 104
- spectrum of the Fourier transform, 116
- spherical maximal function, 475
- spherical average, 476
- spherical coordinates, 591
- spherical Dirichlet kernel, 178
- spherical partial sum, 184
- square Dirichlet kernel, 178
- square function, 421, 458
 - dyadic martingale, 469
- square function of Littlewood–Paley, 421
- square partial sum, 184
- standard kernel, 532
- Stein's interpolation theorem, 41
- stereographic projection, 594
- Stirling's formula, 567
- stopping-time, 97
- stopping-time argument, 356
- strong maximal function, 101
- strong type (p, q) , 33
- sublinear operator, 33, 61
- summation by parts, 599
- support of a distribution, 125
- surface area of the unit sphere \mathbf{S}^{n-1} , 565

- Taylor formula, 607
- tempered distribution, 120
- test function, 119
- tight frame, 496
- tiling of \mathbf{R}^n , 428
- topological group, 18
- torus, 174
- total order of differentiation, 104
- transference of maximal multipliers, 281
- transference of multipliers, 275
- translation
 - of a function, 109
 - of a tempered distribution, 124
- translation operator, 271
- translation-invariant operator, 146
- transpose of an operator, 150
- trigonometric monomial, 177
- trigonometric polynomial, 177
- truncated Hilbert transform, 314
- truncated maximal singular integral, 374
- truncated singular integral, 334, 363

- uncentered Hardy–Littlewood maximal function, 87
- uncentered maximal function with respect to a general measure, 98
- uncentered maximal function with respect to cubes, 99
- uncertainty principle, 118
- uniform boundedness principle, 602
- unitary matrix, 387

- Vandermonde determinant, 165
- variation of a function, 199
- vector-valued
 - extrapolation theorem, 554
 - Hardy–Littlewood maximal inequality, 412, 557
 - inequalities, 555
- vector-valued extension of a linear operator, 390
- vector-valued inequalities, 408, 409, 414
- vector-valued Littlewood–Paley theorem, 426
- vector-valued singular integral, 401
- volume of the unit ball in \mathbf{R}^n , 565

- wave equation, 475
- wavelet, 482
 - of minimally supported frequency, 486
- wavelet transform, 496
- weak L^p , 5
- weak type $(1, 1)$, 88
- weak type (p, q) , 33
- Weierstrass approximation theorem, 32
 - for trigonometric polynomials, 183
- Weierstrass’s theorem, 229
- weight, 499
 - of class A_1 , 502
 - of class A_∞ , 525
 - of class A_p , 503
- weighted estimates
 - for singular integral operators, 540
- Weyl’s theorem, 302
- Whitney decomposition, 609

- Young’s covering lemma, 98
- Young’s inequality, 22, 400
 - for weak type spaces, 23, 73