

Index

Symbols

$[\alpha]$ 114
 $[\cdot]$ 111, 433
 $\langle \alpha \rangle$ 114
 $\langle \cdot \rangle$ 111, 433
 \exists 49, 111, 114, 433
 \forall 48, 111, 114, 433
 \wedge 111, 180, 433
 \leftrightarrow 111
 \rightarrow 111, 180, 433
 \neg 111, 180, 433
 \vee 111, 180, 433
 $?Q$ 76, 429
 $(F)'$ 329
 $(e)'$ 296
 $:$ 76, 429
 F_x^e 591
 $F_x^{c/b}$ 589
 $F_x^{-\infty}$ 612
 $F_x^{e+\varepsilon}$ 617
 F_x^e 595
 $[\cdot]$ 149
 $[\ :=] =$ 204, 368
 $[?]$ 151
 $[\cap]$ 505
 $[\cdot]$ 484
 $[\cup]$ 144
 $[\wedge]$ 161, 231, 336
 $[^d]$ 505
 $[*]$ 155, 216
 $[:=]$ 147, 536
 $[;]$ 153
 $[**]$ 240
 $[\times]$ 505
 $\frac{d}{dt}$ 51, 304, 327
 $\frac{\partial f}{\partial x}(\xi)$ **300**

$\Gamma \vdash_{dGL} P$ **499**
 $\Phi \vdash_{dL} Q$ **180**
 \cdot 537
 \cap 427
 $C^k(D, \mathbb{R}^n)$ **54**
 \cup 76, 429
 $\delta_\alpha(X)$ 474
 \mathbb{R} 195
 $x := *$ 390
 $\exists \text{lin}$ 590
 $\vdash_{dL} P$ **180**, 186, 205
 $\vdash_{dGL} P$ **499**, 500
 $\zeta_\alpha^n(X)$ 461
 ∞ 610
 $\langle ' \rangle$ 487
 $\langle := \rangle$ 486
 $\langle ? \rangle$ 488
 $\langle \cap \rangle$ 505
 $\langle \cdot \rangle$ 157
 $\langle ^d \rangle$ 491
 $\langle * \rangle$ 493
 $\langle ; \rangle$ 490
 $\langle \cup \rangle$ 489
 $\langle \times \rangle$ 505
 \mathcal{S} **45**, 77
 $\Gamma \vdash \Delta$ 177
 ψ **45**
 $BV(P)$ 166
 $BV(\alpha)$ **165**, 166
 $FV(P)$ **165**, 167
 $FV(\alpha)$ **165**, 167
 $FV(\sigma)$ **543**
 $FV(e)$ **165**
 $MBV(\alpha)$ 167
 $V(P)$ 168
 $V(\alpha)$ 168
 $\llbracket P \rrbracket$ 113, 452

$\llbracket \alpha \rrbracket$ 79
 $\llbracket e \rrbracket$ 46
 $\omega \models P$ 48
 $\sigma(\alpha)$ 537
 $\sigma(\phi)$ 537
 $\sigma(\theta)$ 537
 $\zeta_\alpha(X)$ 474
 ε 614
 $\varphi \models x' = f(x) \wedge Q$ 80
 $\varphi \models x' = f(x)$ 80
 $c()$ 533
 $f(e)$ 534
 $p(e)$ 534
 $x' = f(x) \& Q$ 76, 429
 $x' = f(x)$ 76, 429
 $x := e$ 76, 429
 $:=$ 76, 429
 $*$ 76, 429
 \times 428
 d 429
 QE 197, 583
 $i\forall$ 198
 \vdash 177

A

abstraction 70
 admissible *see* substitution, admissible
 advance notice 458
 adversarial 8, 422
 analytic cut *see* cut
 and-operator 52, 123, 443
 Angel **426**
 antecedent **177**
 arity **533**
 assignment **76, 147**
 differential 307, 327
 nondeterministic **390**
 automaton
 hybrid 85, 91
 autonomous 38
 axiom 142–162, 215–217, 301–308,
 327–349, 378–380, 484–494, **527**
 assignment **147, 486, 536**
 composition **153, 490**
 determinacy **484**
 duality **157, 491**
 iteration **155, 216, 493**
 modus ponens
 modal **161**
 of choice **144**
 of game of choice **489**
 solution **149, 487**
 test **151, 488**

 vacuous **162, 231, 536**
 axiomatics 138
 axiomatization 502, 551

B

bi-implication 52, 123, 443
 box-modality 123, 443
 brackets
 semantic 47

C

calculus
 axiomatic *see* axiom
 sequent **179**
 Cauchy-Lipschitz 54
 CEL rule 188
 CER rule 188
 CE rule 188
 challenge **430**
 change
 explicit 52, **308**
 choice
 angelic **426**
 demonic **427**
 nondeterministic **76, 144**
 clash **538, 543**
 closed
 formula *see* formula, closed
 closure
 universal **196**
 competition
 adversarial 437
 analytic 422
 complete **502, 551**
 composition
 sequential **76, 153**
 compositional 138, 298, 455
 conclusion **179**
 condition
 intermediate 125
 winning 433
 conjunction 52, 123, 443
 connectives **174**
 consequence **114**
 consistent **483**
 constant 224, 230
 context **188, 188**
 continued
 to boundary 53
 continuous 8
 evolution *see* differential, equation
 Lipschitz *see* Lipschitz, continuous

contract 97–106
 invariant 107
 contraction 202
 control
 and modeling **12**
 event-triggered **261**, 245–264
 model-predictive **273**
 time-triggered **280**, 267–282
 convention
 naming 44
 counterexample **121**
 CPS 1
 cut **183**
 differential *see* differential, cut

D

dA rule 380
 DC axiom 349
 dC rule 346
 DE axiom 307, 327
 De Morgan 586
 degree **611**
 Demon **426**
 derivation 302
 operator **317**
 derivative
 partial **300**
 descriptive power 39
 design
 iterative 69
 determined **483**
 DG axiom 379
 dG rule 380
 dGL 433
 DI axiom 305, 340
 dI rule 309, 331, 340
 diamond-modality 123, 443
 differential 295, 300, 304, 327
 algebra 317
 auxiliaries **380**
 axiomatization 388
 constant 317
 cut 346, 349
 effect 307, 327
 equation **32**, 28–58, **76**, 80, 148–151,
 287–415
 dual **428**
 field of quotients **317**
 formula 329
 ghost **379**, 380
 indeterminates 317
 invariant 331, 340
 polynomial algebra **317**

 structure 404
 symbol 297, 339
 term 296, 300
 weakening 327
 discrete 8
 disjunction 52, 123, 443
 distributed 8
 dL **111**
 domain **32**
 evolution **40**, **76**, **429**
 reduction 503
 of substitution **537**
 doubt
 Cartesian 123
 draw **482**
 DW axiom 327
 dW rule 328
 dynamics
 adversarial **422**
 continuous 39–52
 discrete 66

E

equation
 differential *see* differential, equation
 equilibrium **385**
 equivalence
 contextual 188, 551
 equivalent 52, 123, 443
 event 251
 evolution
 continuous *see* differential, equation
 exchange 201
 existence
 of solution 53
 existential quantifier *see* quantifier,
 existential
 exists 52
 expression
 square root **592**
 expressive
 differentially **502**, 551

F

filibuster 441
 fixpoint **472**
 for all 52
 form
 differential 289
 game **429**
 formula
 dGL **433**

dL 111
 atomic **183**
 closed 197, 584
 differential game logic **433**
 differential dynamic logic **111**
 with interpretation 534
 FOL **43**
 ground 197, 584
 modal **110, 433**
 principal 183
 FP[×] rule 506
 FP rule 494

G

Gödel 162, 218
 G rule 162, 218
 game 424–432
 atomic **429**
 choice **430**
 compound **430**
 continuous evolution **429**
 demonic choice 427
 demonic repetition 428
 deterministic assignment **429**
 dual **430**
 dual test **430**
 hybrid 8, **429**
 repeated **430**
 sequential **430**
 test **430**
 generator 316
 George Box 561
 ghost
 arithmetic 390
 axiomatic **387**
 differential *see* differential, ghost
 differential-algebraic 393
 discrete 367
 substitute **383**
 graph **53**
 ground
 formula *see* formula, ground

H

HG 429
 hide 202
 Hoare
 triple **117**
 homomorphism **542**
 HP 76
 hybrid *see* system, hybrid

I

I axiom 217
 IA rule 367
 ideal
 vanishing **600**
 implication 52, 123, 443
 implies-operator 52, 123, 443
 ind[×] rule 506
 ind rule 219
 inductive **221**
 infinitesimal **614**
 inflation 473
 initial value
 problem 32
 symbolic 34, 149
 instantiation 200
 intermediate
 condition 125
 state 124
 interpretation **535**
 adjoint **546**
 invariant **221**
 differential 305, 331, 340
 function **319**
 term 305

K

kernel **48**
 K axiom 161

L

Leibniz
 rule **317**
 Lie-derivative 319, **339**
 Lipschitz **54**
 local **54**
 logic
 differential dynamic 107–115
 with interpretation 533–536
 differential game 428–439
 first-order 42–50, 580
 loop
 invariant 221
 rule 221

M

M rule 163, 220
 M[·] rule 163, 220
 [·]-modality 123, 443
 ⟨·⟩-modality 123, 443

ModelPlex 557–574
 model checking
 bounded 235
 monitor **569**
 controller **563**
 model **564**
 runtime **569**
 monotone **456, 481**
 MR rule 237
 multi-firing 259
 M rule 485
 M[.] rule 485

N

negation 52, 123, 443
 nilpotent **56**
 nondeterministic 8, **70**
 not-operator 52, 123, 443

O

objective 433
 Occam 201
 ODE *see* differential, equation
 operator
 duality **429**
 modal **110**
 precedence 77, **113**
 or-operator 52, 123, 443
 ordinal **465**
 arithmetic 468
 limit **465, 467**
 successor **465, 467**
 transfinite **466**

P

parameter
 constant 224, 230
 symbolic 119
 Peano 53
 Picard-Lindelöf 54
 postcondition **110**
 powerset **81, 115**
 pre-fixpoint **470**
 precedence
 operator **113**
 premise **179**
 program
 atomic **76**
 compound **76**
 continuous 42
 hybrid **76, 65–80, 534**

proof **179**
 irrelevance 255
 propositional
 rule *see* rule, propositional
 proved **180, 499**

Q

QE (quantifier elimination) 197, 583
 quantifier
 elimination **197, 583**
 equational 591
 linear 589
 quadratic 618
 existential 52, 123, 443
 universal 52, 123, 443
 Quantum 98
 quotient
 rule **317**

R

reaction 273
 reals 43
 extended 613
 nonstandard 618
 region
 winning **452, 474**
 iterated **461**
 repetition
 angelic **426**
 demonic **428**
 nondeterministic **76, 155, 216**
 response 246
 rule 162–164, 180–205, 217–223, 327–349,
 367, 378–380, 494–495, 538, 552
 arithmetic 195
 axiomatic **552**
 derived **192**
 fixpoint **494**
 Gödel **162, 218**
 loop 221
 loop **494**
 monotonicity **163, 220, 485**
 product **317**
 proof 179
 propositional 175–193

S

satisfaction **48, 114, 453**
 satisfiable 49
 schema
 axiom **527**

- rule 551
 - semantics **113, 138, 535**
 - denotational **298**
 - of dGL 452
 - of dL 112
 - of differentials 300
 - of differential symbols 297
 - of FOL 47
 - of games 474
 - of HPs 77
 - of terms 45
 - operational 439
 - static **164**
 - sequent **177, 179**
 - set
 - closed **411**
 - open **411**
 - semialgebraic **625**
 - skills
 - CPS **14**
 - Skolem
 - function 190
 - solution **32, 51, 80, 149, 369**
 - soundness
 - local **552**
 - miracle 500
 - of axioms **146, 159, 548**
 - of calculus **186, 205**
 - of dL 159
 - of proof rule **179**
 - of dGL 500
 - of US 547, 552
 - state **45, 77, 297, 451**
 - modification **49**
 - set of **45, 77, 451**
 - stochastic 8
 - strategy **441**
 - winning **433, 433, 441, 474**
 - subgame 429
 - substitution
 - admissible 148, **543**
 - uniform **536, 542, 536–547**
 - of rule 552
 - virtual 590–599, 609–624
 - infinitesimal **615**
 - infinity **610**
 - square root **595**
 - succedent **177**
 - symbol **531**
 - constant **533**
 - differential **295**
 - function **533**
 - predicate **533**
 - program constant **533**
 - syntax 138
 - of dGL 433
 - of dL 110
 - of FOL 43
 - of HGs 429
 - of HPs 75
 - of terms 42
 - system
 - cyber-physical 1
 - hybrid **6, 8**
 - distributed 8
 - stochastic 8
 - multi-dynamical **8, 422**
- T**
- tangent 300
 - Tarski 197, 584
 - Tarski-Seidenberg 625
 - term **42, 296, 534**
 - atomic **43**
 - compound **43**
 - differential **295, 300**
 - test **76, 151, 429**
 - dual **428**
 - poor **503**
 - rich **503**
 - theory
 - proof **399, 502, 551**
 - thinking
 - computational **13**
 - time 38, 82, 271
 - divergence of **278**
 - topology 411
 - transfinite 473
 - transition
 - semantics
 - dL 78
 - trinity 138
 - logical **138**
- U**
- uncertain 8
 - uniform substitution *see* substitution, uniform
 - uniqueness
 - of solution 54
 - universal quantifier *see* quantifier, universal
 - unsatisfiable 49
 - US rule 538, 547
- V**
- V axiom 162, 231
 - vacuous 162, 231, 536

valid 49, **114**, **453**, **535**

value

of term **46**, **535**

of term

differential 300

variable 42

bound **164**, **166**

differential *see* symbol, differential

free **164**, **167**

must-bound **167**

variant **516**

variety

affine **600**

velocity

limit **385**

V axiom 536

W

ω -semantics 461

weakening 202

differential *see* differential, weakening

Z

Zeno **278**

zero-sum **482**

Operators of Differential Dynamic Logic (dL)

dL	Operator	Meaning
$e \geq \tilde{e}$	greater or equals	true if value of e greater-or-equal to \tilde{e}
$\neg P$	negation / not	true if P is false
$P \wedge Q$	conjunction / and	true if both P and Q are true
$P \vee Q$	disjunction / or	true if P is true or if Q is true
$P \rightarrow Q$	implication / implies	true if P is false or Q is true
$P \leftrightarrow Q$	bi-implication / equivalent	true if P and Q are both true or both false
$\forall x P$	universal quantifier / for all	true if P is true for all values of variable x
$\exists x P$	existential quantifier / exist	true if P is true for some value of variable x
$[\alpha]P$	$[\cdot]$ modality / box	true if P is true after all runs of HP α
$\langle \alpha \rangle P$	$\langle \cdot \rangle$ modality / diamond	true if P is true after some run of HP α

Statements and effects of Hybrid Programs (HPs)

HP Notation	Operation	Effect
$x := e$	discrete assignment	assigns current value of term e to variable x
$x := *$	nondet. assignment	assigns any real value to variable x
$x' = f(x) \ \& \ Q$	continuous evolution	follow differential equation $x' = f(x)$ within evolution domain Q for any duration
$?Q$	state test / check	test first-order formula Q at current state
$\alpha; \beta$	seq. composition	HP β starts after HP α finishes
$\alpha \cup \beta$	nondet. choice	choice between alternatives HP α or HP β
α^*	nondet. repetition	repeats HP α any $n \in \mathbb{N}$ times

Semantics of dL formula P is the set of states $\llbracket P \rrbracket \subseteq \mathcal{S}$ in which it is true

$$\begin{aligned} \llbracket e \geq \tilde{e} \rrbracket &= \{ \omega \in \mathcal{S} : \omega[e] \geq \omega[\tilde{e}] \} \\ \llbracket P \wedge Q \rrbracket &= \llbracket P \rrbracket \cap \llbracket Q \rrbracket \\ \llbracket P \vee Q \rrbracket &= \llbracket P \rrbracket \cup \llbracket Q \rrbracket \\ \llbracket \neg P \rrbracket &= \llbracket P \rrbracket^c = \mathcal{S} \setminus \llbracket P \rrbracket \\ \llbracket \langle \alpha \rangle P \rrbracket &= \llbracket \alpha \rrbracket \circ \llbracket P \rrbracket = \{ \omega : \nu \in \llbracket P \rrbracket \text{ for some state } \nu \text{ such that } (\omega, \nu) \in \llbracket \alpha \rrbracket \} \\ \llbracket [\alpha] P \rrbracket &= \llbracket \neg \langle \alpha \rangle \neg P \rrbracket = \{ \omega : \nu \in \llbracket P \rrbracket \text{ for all states } \nu \text{ such that } (\omega, \nu) \in \llbracket \alpha \rrbracket \} \\ \llbracket \exists x P \rrbracket &= \{ \omega : \nu \in \llbracket P \rrbracket \text{ for some state } \nu \text{ that agrees with } \omega \text{ except on } x \} \\ \llbracket \forall x P \rrbracket &= \{ \omega : \nu \in \llbracket P \rrbracket \text{ for all states } \nu \text{ that agree with } \omega \text{ except on } x \} \end{aligned}$$

Semantics of HP α is relation $\llbracket \alpha \rrbracket \subseteq \mathcal{S} \times \mathcal{S}$ between initial and final states

$$\begin{aligned} \llbracket x := e \rrbracket &= \{ (\omega, \nu) : \nu = \omega \text{ except that } \nu[x] = \omega[e] \} \\ \llbracket ?Q \rrbracket &= \{ (\omega, \omega) : \omega \in \llbracket Q \rrbracket \} \\ \llbracket x' = f(x) \ \& \ Q \rrbracket &= \{ (\omega, \nu) : \varphi(0) = \omega \text{ except at } x' \text{ and } \varphi(r) = \nu \text{ for a solution } \varphi: [0, r] \rightarrow \mathcal{S} \text{ of any duration } r \text{ satisfying } \varphi \models x' = f(x) \wedge Q \} \\ \llbracket \alpha \cup \beta \rrbracket &= \llbracket \alpha \rrbracket \cup \llbracket \beta \rrbracket \\ \llbracket \alpha; \beta \rrbracket &= \llbracket \alpha \rrbracket \circ \llbracket \beta \rrbracket = \{ (\omega, \nu) : (\omega, \mu) \in \llbracket \alpha \rrbracket, (\mu, \nu) \in \llbracket \beta \rrbracket \} \\ \llbracket \alpha^* \rrbracket &= \llbracket \alpha \rrbracket^* = \bigcup_{n \in \mathbb{N}} \llbracket \alpha^n \rrbracket \text{ with } \alpha^{n+1} \equiv \alpha^n; \alpha \text{ and } \alpha^0 \equiv \text{true} \end{aligned}$$

Axiomatization (dL)

$$\langle \cdot \rangle \quad \langle \alpha \rangle P \leftrightarrow \neg[\alpha]\neg P$$

$$[:=] \quad [x:=e]p(x) \leftrightarrow p(e)$$

$$[?] \quad [?Q]P \leftrightarrow (Q \rightarrow P)$$

$$['] \quad [x' = f(x)]p(x) \leftrightarrow \forall t \geq 0 [x := y(t)]p(x) \quad (y'(t) = f(y))$$

$$[\cup] \quad [\alpha \cup \beta]P \leftrightarrow [\alpha]P \wedge [\beta]P$$

$$[:] \quad [\alpha; \beta]P \leftrightarrow [\alpha][\beta]P$$

$$[*] \quad [\alpha^*]P \leftrightarrow P \wedge [\alpha][\alpha^*]P$$

$$\text{K} \quad [\alpha](P \rightarrow Q) \rightarrow ([\alpha]P \rightarrow [\alpha]Q)$$

$$\text{I} \quad [\alpha^*]P \leftrightarrow P \wedge [\alpha^*](P \rightarrow [\alpha]P)$$

$$\forall p \rightarrow [\alpha]p \quad (FV(p) \cap BV(\alpha) = \emptyset)$$

$$\text{M}[\cdot] \quad \frac{P \rightarrow Q}{[\alpha]P \rightarrow [\alpha]Q}$$

$$\text{G} \quad \frac{P}{[\alpha]P}$$

Differential equation axioms

$$\text{DW} \quad [x' = f(x) \& Q]P \leftrightarrow [x' = f(x) \& Q](Q \rightarrow P)$$

$$\text{DI} \quad ([x' = f(x) \& Q]P \leftrightarrow [?Q]P) \leftarrow (Q \rightarrow [x' = f(x) \& Q](P)')$$

$$\text{DC} \quad ([x' = f(x) \& Q]P \leftrightarrow [x' = f(x) \& Q \wedge C]P) \leftarrow [x' = f(x) \& Q]C$$

$$\text{DE} \quad [x' = f(x) \& Q]P \leftrightarrow [x' = f(x) \& Q][x' := f(x)]P$$

$$\text{DG} \quad [x' = f(x) \& Q]P \leftrightarrow \exists y [x' = f(x), y' = a(x) \cdot y + b(x) \& Q]P$$

$$+' \quad (e+k)' = (e)' + (k)'$$

$$\cdot' \quad (e \cdot k)' = (e)' \cdot k + e \cdot (k)'$$

$$c' \quad (c())' = 0 \quad (\text{for numbers or constants } c())$$

$$x' \quad (x)' = x' \quad (\text{for variable } x \in \mathcal{V})$$

Differential equation proof rules

$$\text{dW} \quad \frac{Q \vdash P}{\Gamma \vdash [x' = f(x) \& Q]P, \Delta} \quad \text{dI} \quad \frac{Q \vdash [x' := f(x)](F)'}{F \vdash [x' = f(x) \& Q]F}$$

$$\text{dC} \quad \frac{\Gamma \vdash [x' = f(x) \& Q]C, \Delta \quad \Gamma \vdash [x' = f(x) \& (Q \wedge C)]P, \Delta}{\Gamma \vdash [x' = f(x) \& Q]P, \Delta}$$

Sequent calculus proof rules

$$\begin{array}{l}
\text{\textasciitilde}R \frac{\Gamma, P \vdash \Delta}{\Gamma \vdash \text{\textasciitilde}P, \Delta} \quad \wedge R \frac{\Gamma \vdash P, \Delta \quad \Gamma \vdash Q, \Delta}{\Gamma \vdash P \wedge Q, \Delta} \quad \vee R \frac{\Gamma \vdash P, Q, \Delta}{\Gamma \vdash P \vee Q, \Delta} \\
\text{\textasciitilde}L \frac{\Gamma \vdash P, \Delta}{\Gamma, \text{\textasciitilde}P \vdash \Delta} \quad \wedge L \frac{\Gamma, P, Q \vdash \Delta}{\Gamma, P \wedge Q \vdash \Delta} \quad \vee L \frac{\Gamma, P \vdash \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \vee Q \vdash \Delta} \\
\text{\textasciitilde}\rightarrow R \frac{\Gamma, P \vdash Q, \Delta}{\Gamma \vdash P \rightarrow Q, \Delta} \quad \text{id} \frac{}{\Gamma, P \vdash P, \Delta} \quad \text{WR} \frac{\Gamma \vdash \Delta}{\Gamma \vdash P, \Delta} \\
\text{\textasciitilde}\rightarrow L \frac{\Gamma \vdash P, \Delta \quad \Gamma, Q \vdash \Delta}{\Gamma, P \rightarrow Q \vdash \Delta} \quad \text{cut} \frac{\Gamma \vdash C, \Delta \quad \Gamma, C \vdash \Delta}{\Gamma \vdash \Delta} \quad \text{WL} \frac{\Gamma \vdash \Delta}{\Gamma, P \vdash \Delta} \\
\forall R \frac{\Gamma \vdash p(y), \Delta}{\Gamma \vdash \forall x p(x), \Delta} \quad (y \notin \Gamma, \Delta, \forall x p(x)) \quad \exists R \frac{\Gamma \vdash p(e), \Delta}{\Gamma \vdash \exists x p(x), \Delta} \quad (\text{arbitrary term } e) \\
\forall L \frac{\Gamma, p(e) \vdash \Delta}{\Gamma, \forall x p(x) \vdash \Delta} \quad (\text{arbitrary term } e) \quad \exists L \frac{\Gamma, p(y) \vdash \Delta}{\Gamma, \exists x p(x) \vdash \Delta} \quad (y \notin \Gamma, \Delta, \exists x p(x)) \\
\text{CER} \frac{\Gamma \vdash C(Q), \Delta \quad \vdash P \leftrightarrow Q}{\Gamma \vdash C(P), \Delta} =R \frac{\Gamma, x = e \vdash p(e), \Delta}{\Gamma, x = e \vdash p(x), \Delta} \\
\text{CEL} \frac{\Gamma, C(Q) \vdash \Delta \quad \vdash P \leftrightarrow Q}{\Gamma, C(P) \vdash \Delta} =L \frac{\Gamma, x = e, p(e) \vdash \Delta}{\Gamma, x = e, p(x) \vdash \Delta}
\end{array}$$

Derived axioms and derived rules

$$\begin{array}{l}
\wedge' (P \wedge Q)' \leftrightarrow (P)' \wedge (Q)' \\
\vee' (P \vee Q)' \leftrightarrow (P)' \vee (Q)' \\
\llbracket \wedge [\alpha](P \wedge Q) \leftrightarrow [\alpha]P \wedge [\alpha]Q \\
\llbracket * [\alpha^*]P \leftrightarrow P \wedge [\alpha^*][\alpha]P \\
\llbracket ** [\alpha^*; \alpha^*]P \leftrightarrow [\alpha^*]P \\
[:=] = \frac{\Gamma, y = e \vdash p(y), \Delta}{\Gamma \vdash [x := e]p(x), \Delta} \quad (y \text{ new}) \\
\text{iG} \frac{\Gamma \vdash [y := e]p, \Delta}{\Gamma \vdash p, \Delta} \quad (y \text{ new}) \\
\text{dG} \frac{\Gamma \vdash \exists y [x' = f(x), y' = a(x) \cdot y + b(x) \& Q]P, \Delta}{\Gamma \vdash [x' = f(x) \& Q]P, \Delta} \\
\text{dA} \frac{\vdash J \leftrightarrow \exists y G \quad G \vdash [x' = f(x), y' = a(x) \cdot y + b(x) \& Q]G}{J \vdash [x' = f(x) \& Q]J}
\end{array}$$