

# Chapter 15

## Trigonometric Functions

### Statement of the Problem

*Mathematica* can calculate Cos and Sin for many arguments equal to  $\pi$  times rational numbers. For example,

**In[1] := Table[Cos[Pi/n], {n, 1, 12}]**

Out[1] =  $\left\{ -1, 0, \frac{1}{2}, \frac{1}{\sqrt{2}}, \frac{1}{4} (1 + \sqrt{5}), \frac{\sqrt{3}}{2}, \text{Cos} \left[ \frac{\pi}{7} \right], \text{Cos} \left[ \frac{\pi}{8} \right], \text{Cos} \left[ \frac{\pi}{9} \right], \right.$   
 $\left. \sqrt{\frac{5}{8} + \frac{\sqrt{5}}{8}}, \text{Cos} \left[ \frac{\pi}{11} \right], \frac{1 + \sqrt{3}}{2\sqrt{2}} \right\}$

But it does not apply the half-angle formulas in all possible cases. We'll write our own cos function which does this; then

**In[2] := sin[x\_] := cos[x - Pi/2]**

### Simple Cases

These cases should be considered separately because *Mathematica* does not treat them as rational numbers times  $\pi$ .

**In[3] := cos[0] = 1; cos[Pi] = -1;**

**In[4] := cos[n\_Integer \* Pi] := (-1)^n**

### General Case

**In[5] := cos[r\_Rational \* Pi] := Which[r < 0, cos[-r \* Pi],  
 r > 2, cos[FractionalPart[r/2] \* 2 \* Pi], r > 1, cos[(2 - r) \* Pi],  
 r > 1/2, -cos[(1 - r) \* Pi],**

**EvenQ[Denominator[r]], Simplify[Sqrt[(1 + cos[2 \* r \* Pi])/2]],  
True, Cos[r \* Pi]]**

For example,

**In[6] := {cos[Pi/32], cos[-65 \* Pi/32], cos[3 \* Pi/32], cos[33 \* Pi/32]}**

$$\text{Out[6]} = \left\{ \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}}, \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}}, \right. \\ \left. \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 - \sqrt{2}}}}, -\frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}} \right\}$$

**In[7] := {cos[Pi/48], cos[5 \* Pi/48], cos[7 \* Pi/48]}**

$$\text{Out[7]} = \left\{ \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{3}}}}, \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 - \sqrt{3}}}}, \right. \\ \left. \frac{1}{2} \sqrt{2 + \sqrt{2 - \sqrt{2 - \sqrt{3}}}} \right\}$$

**In[8] := {cos[Pi/40], cos[3 \* Pi/40]}**

$$\text{Out[8]} = \left\{ \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{\frac{1}{2} (5 + \sqrt{5})}}}, \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{\frac{1}{2} (5 - \sqrt{5})}}} \right\}$$

### Check

**In[9] := check[d\_, n\_] := Module[{e = 0, x, ex},  
Do[x = i/d \* Pi; ex = Abs[N[cos[x] - Cos[x]]]; If[ex > e, e = ex,  
{i, -d \* n, d \* n}];**

**e]**

**In[10] := {check[128, 5], check[192, 5], check[320, 5]}**

$$\text{Out[10]} = \left\{ 2.220446049250313 \times 10^{-16}, 6.106226635438361 \times 10^{-16}, \right. \\ \left. 1.861358289723114 \times 10^{-15} \right\}$$