

# Chapter 16

## Cerebellar Nucleus Development

Hong-Ting Prekop and Richard J.T. Wingate

**Abstract** The cerebellar nuclei are the final output structures of the cerebellum, integrating extra-cerebellar inputs from climbing fibres and mossy fibres with inhibitory signals from Purkinje cells. As in overlying cortical layers, cerebellar nuclei contain a mixture of glutamatergic and GABAergic neurons, which develop from the rhombic lip and ventricular zone, respectively.

**Keywords** Atoh (atonal) 1 • Ptf1a • Climbing fibre • Mossy fibre • GABAergic neuron • Glutamatergic neurons • Neuronal migration • Rhombic lip


The cerebellar nuclei (CN) are the final output structures of the cerebellum, integrating inputs from the forebrain, brainstem and spinal cord with the cerebellar cortical output from Purkinje cells. In contrast to the remarkably evolutionarily conserved connectivity between granule cells and Purkinje cells in the cerebellar cortex, the size, foliation and number of CN varies between animals (one in amphibians, two in reptiles and birds, three to five in mammals) (Nieuwenhuys et al. 1998). In humans, there are four nuclei: the medial (fastigial), anterior and posterior interposed, and lateral (dentate) nuclei.

Much of what is known concerning CN cell morphologies and neuronal circuitry comes from Golgi and Nissl preparations of the mammalian lateral nucleus observed by light and electron microscopy (Chan-Palay 1977). Based on neurotransmitter content and neuronal connectivity, four main cell types have been identified: large excitatory glutamatergic projection neurons, inhibitory GABAergic projection neurons and local, inhibitory GABAergic and glycinergic interneurons (Fig. 16.1) (Batini et al. 1992; Chen and Hillman 1993; Fredette et al. 1992).

CN receive inhibitory projections from the Purkinje cells from the overlying cerebellar cortex in a broadly topographic manner. The lateral nuclei are innervated by the lateral cerebellum and medial nuclei by the medial vermis (Voogd and Glickstein 1998). CN also integrate collateral inputs from axons projecting from the

---

H.-T. Prekop (✉) • R.J.T. Wingate  
Department of Developmental Neurobiology, King's College London,  
4th floor New Hunts House, Guys Campus, London SE1 1UL, UK  
e-mail: [hong-ting.kwok@kcl.ac.uk](mailto:hong-ting.kwok@kcl.ac.uk)




**Fig. 16.1** Connections and circuitry of the human lateral nucleus. *Left:* The lateral CN receives input from Purkinje Cells (PC) and innervates the thalamus, which in turn modulates cortical activity. The nucleus also receives climbing fibres (CF) collateral input from the inferior olive (IO), and mossy fibres (MF) collaterals, originating principally from the pons. Climbing and mossy fibres terminate on PC and granule cells (GC) in the cerebellar cortex, respectively. The CN also contains GABAergic projection neurons that send inhibitory signals to the IO. *Right:* Within the nucleus, each PC axon fans out into a cone that innervates a number of excitatory and GABAergic projection neurons. Small GABAergic and glycinergic interneurons provide local inhibition. Inputs from MF and CF collaterals run perpendicular to the afferent Purkinje cell axons (Adapted from Chan-Palay 1977)

pontine nucleus and inferior olive to the cerebellar cortex (Fig. 16.1). The output of each CN is then directed to different central neural systems as determined by the pattern of their efferent connections (Larsell and Jansen 1972). Of these, the connection from the lateral nucleus to ventrolateral thalamus is a uniquely mammalian adaptation and completes a closed-loop cortico-pontine-cerebellar relay circuit (Kelly and Strick 2003) that is heavily implicated in modulating higher cognitive function in humans (Schmahmann 2010).

## 16.1 Concepts of CN Development Have Changed Markedly in Recent Years

Cell types of the cerebellum arise from two germinal regions within the most anterior neuromere of embryonic hindbrain, rhombomere 1: the ventricular zone (VZ) and the rhombic lip (RL). The VZ is a neuroepithelial zone that lines the dorsolateral part of the fourth ventricle, while the rhombic lip comprises the interface of this neuroepithelium with the roof plate of the fourth ventricle (Wingate 2001). Until the last decade, it was thought that all CN neurons originate from the VZ then migrate radially into the white matter (Altman and Bayer 1985a, b; Goldowitz and Hamre 1998). It is now known that CN neurons of different neurotransmitter types are born from both the RL and VZ.



**Fig. 16.2** Cerebellar nucleus neurons have a dual origin. Different neuronal subtypes in the cerebellum develop from separate germinal regions. Precursors continue to migrate, proliferate, differentiate and mature postnatally. The CN glutamatergic cells form first at the RL and migrate via the nuclear transitory zone (NTZ). The RL also gives rise to granule cell precursors, which form the external granule layer (EGL) that later migrates in to form the inner granule layer (IGL). All cortical GABAergic cells originate from the VZ, migrate into the intermediate zone (IZ) and finally to the molecular (ML) and granule (GL) cell layers

Research by classical birth dating and genetic fate mapping has shown that all cerebellar excitatory neurons are derived from RL progenitors, specified by the expression of *Atoh1*, while inhibitory neurons arise from the VZ, where progenitors are specified by the early expression of *Ptfla* (Hoshino et al. 2005). The bHLH proteins, *Ptfla* and *Atoh1*, have both been shown to be necessary (Machold and Fishell 2005; Wang et al. 2005) and sufficient (Yamada et al. 2014) for the production of all GABAergic and glutamatergic neurons in the cerebellum, respectively (Fig. 16.2). The following description exemplifies CN development using the embryonic mouse model over its 21 days of gestation.

### 16.1.1 Glutamatergic Neurons Are Born at the Rhombic Lip

*Atoh1* expressing progenitor cells from the RL produce glutamatergic CN projection neurons between embryonic day (e)10–e12.5 prior to making granule cell precursors that populate the external granule layer (EGL) (Machold and Fishell 2005; Wang et al. 2005). From e12.5–e14.5, the CN cells migrate from the RL across the dorsal surface of rhombomere 1 via the subplial rhombic lip migratory stream (RLS), then congregate at the nuclear transitory zone (NTZ) at the boundary of the cerebellar anlage (Fig. 16.2). The NTZ is thought to be a transient differentiation zone (Altman and Bayer 1985a), where nuclear neurons are defined by specific, temporally restricted, developmental transcription factor profiles (Fink et al. 2006).

The CN are born in a lateral to medial sequence subsequent to the first-born RL derivatives, which become extra-cerebellar neurons (Machold and Fishell 2005). Like these extra-cerebellar neurons, nuclear cells in the lateral nucleus of mammals express the LIM-homeodomain gene *Lhx9* (Wang et al. 2005; Green and Wingate 2014). Both these early populations project to the thalamus suggesting a role for *Lhx9* in specifying axonal projection (Green and Wingate 2014). Subsequently, RL-derived projection neurons of the interposed and medial nuclei are defined by

their expression of *Tbr2* and *Tbr1*, respectively (Fink et al. 2006; Engelkamp et al. 1999; Landsberg et al. 2005) and extend axons to various hindbrain, midbrain and ventral diencephalic targets.

From e14.5–e16.5, CN cells in the NTZ descend into the white matter. It is unclear whether this is due to active migration towards the VZ (Altman and Bayer 1985a) or displacement by gross morphogenic changes to cerebellar shape as granule cell precursors in the EGL proliferate to produce the most abundant neuronal population in the brain.

### ***16.1.2 GABAergic Neurons Are Derived from the Ventricular Zone***

Fate-mapping studies indicate that GABAergic CN cells are derived from *Ptf1a*-positive precursors in the VZ in two phases (Hoshino et al. 2005). First, GABAergic neurons that project long axons from the CN to the inferior olive (Mugnaini and Oertel 1985; Ruigrok 1997) are born within a distinct temporal window alongside Purkinje cells (e10.5–e12.5), both characterised by the expression of *Olig2*. Very little is known about their subsequent temporal expression profiles or migration: for example, it is unclear whether they accumulate alongside excitatory projection neurons in the NTZ before descending to their destination.

From e13.5, *Olig2* is down regulated and subsequent populations of GABAergic neurons express *Gsx1* (Seto et al. 2014) and *Pax2* (Maricich and Herrup 1999; Weisheit et al. 2006). *Pax2*-positive precursors proliferate within the white matter through to P15 and migrate radially to sequentially form various GABAergic interneuron populations: first the CN interneurons, then Golgi cells of the granule cell layer and finally basket and stellate cells of the molecular layer (Leto et al. 2006). A growing body of evidence shows that specification is controlled post-mitotically by factors in the local microenvironment (Leto et al. 2009; Grimaldi et al. 2009; Zordan et al. 2008), although their identity, and the contribution, if any, of intrinsic cues are still largely undefined.

## **16.2 Future Studies Will Need to Address Fine-Grain Patterning of Different Nuclei**

Our current understanding outlines basic principles of CN development in terms of progenitor zones, temporal patterning and the function of a few key transcription factors. Recent studies have illustrated how the discovery of new molecular and genetic markers has allowed fate mapping of distinct cell types. Detailed studies of cell organisation within the lateral nucleus have revealed intricate cell arrangements and alignment of projections along a polarized axis within the nucleus (Chan-Palay 1977). How neuroblasts migrate, differentiate, and successfully form functional

circuits are important open questions. The identity of cues that shape CN circuits will be important targets for future research. This will also help in assessing the impact of CN dysgenesis on a broad spectrum of cerebellar disorders that can produce both classical motor symptoms and an emerging range of cognitive effects in syndromes such as Autistic Spectrum Disorder and Joubert Syndrome (Schmahmann 2010; Wang et al. 2014; Holroyd et al. 1991).

## References

- Altman J, Bayer SA (1985a) Embryonic development of the rat cerebellum. II. Translocation and regional distribution of the deep neurons. *J Comp Neurol* 231:27–41
- Altman J, Bayer SA (1985b) Embryonic development of the rat cerebellum. III. Regional differences in the time of origin, migration, and settling of Purkinje cells. *J Comp Neurol* 231:42–65
- Batini C, Compoin C, Buisseret-Delmas C, Daniel H, Guegan M (1992) Cerebellar nuclei and the nucleocortical projections in the rat: retrograde tracing coupled to GABA and glutamate immunohistochemistry. *J Comp Neurol* 315:74–84
- Chan-Palay V (1977) *Cerebellar dentate nucleus: organization, cytology and transmitters*. Springer, Berlin
- Chen S, Hillman DE (1993) Colocalization of neurotransmitters in the deep cerebellar nuclei. *J Neurocytol* 22:81–91
- Engelkamp D, Rashbass P, Seawright A, van Heyningen V (1999) Role of Pax6 in development of the cerebellar system. *Development* 126:3585–3596
- Fink AJ, Englund C, Daza RA, Pham D, Lau C, Nivison M, Kowalczyk T, Hevner RF (2006) Development of the deep cerebellar nuclei: transcription factors and cell migration from the rhombic lip. *J Neurosci* 26:3066–3076
- Fredette BJ, Adams JC, Mugnaini E (1992) GABAergic neurons in the mammalian inferior olive and ventral medulla detected by glutamate decarboxylase immunocytochemistry. *J Comp Neurol* 321:501–514
- Goldowitz D, Hamre K (1998) The cells and molecules that make a cerebellum. *Trends Neurosci* 21:375–382
- Green MJ, Wingate RJ (2014) Developmental origins of diversity in cerebellar output nuclei. *Neural Dev* 9:1
- Grimaldi P, Parras C, Guillemot F, Rossi F, Wassef M (2009) Origins and control of the differentiation of inhibitory interneurons and glia in the cerebellum. *Dev Biol* 328:422–433
- Holroyd S, Reiss AL, Bryan RN (1991) Autistic features in Joubert syndrome: a genetic disorder with agenesis of the cerebellar vermis. *Biol Psychiatry* 29:287–294
- Hoshino M, Nakamura S, Mori K, Kawauchi T, Terao M, Nishimura YV, Fukuda A, Fuse T, Matsuo N, Sone M et al (2005) Ptf1a, a bHLH transcriptional gene, defines GABAergic neuronal fates in cerebellum. *Neuron* 47:201–213
- Kelly RM, Strick PL (2003) Cerebellar loops with motor cortex and prefrontal cortex of a nonhuman primate. *J Neurosci* 23:8432–8444
- Landsberg RL, Awatramani RB, Hunter NL, Farago AF, Dipietrantonio HJ, Rodriguez CI, Dymecki SM (2005) Hindbrain rhombic lip is comprised of discrete progenitor cell populations allocated by pax6. *Neuron* 48:933–947
- Larsell O, Jansen J (1972) *The comparative anatomy and histology of the cerebellum: the human cerebellum, cerebellar connections, and cerebellar cortex*. University of Minnesota, Minnesota
- Leto K, Carletti B, Williams IM, Magrassi L, Rossi F (2006) Different types of cerebellar GABAergic interneurons originate from a common pool of multipotent progenitor cells. *J Neurosci* 26:11682–11694

- Leto K, Bartolini A, Yanagawa Y, Obata K, Magrassi L, Schilling K, Rossi F (2009) Laminar fate and phenotype specification of cerebellar GABAergic interneurons. *J Neurosci* 29:7079–7091
- Machold R, Fishell G (2005) Math1 is expressed in temporally discrete pools of cerebellar rhombic-lip neural progenitors. *Neuron* 48:17–24
- Maricich SM, Herrup K (1999) Pax-2 expression defines a subset of GABAergic interneurons and their precursors in the developing murine cerebellum. *J Neurobiol* 41:281–294
- Mugnaini E, Oertel W (1985) An atlas of the distribution of GABAergic neurons and terminals in the rat CNS as revealed by GAD immunohistochemistry. In: *Handbook of chemical neuroanatomy*, vol 4. Elsevier, Amsterdam, pp 436–622
- Nieuwenhuys R, ten Donkelaar HJ, Nicholson C (eds) (1998) *The central nervous system of vertebrates*. Springer, Berlin
- Ruigrok TJ (1997) Cerebellar nuclei: the olivary connection. *Prog Brain Res* 114:167–192
- Schmahmann JD (2010) The role of the cerebellum in cognition and emotion: personal reflections since 1982 on the dysmetria of thought hypothesis, and its historical evolution from theory to therapy. *Neuropsychol Rev* 20:236–260
- Seto Y, Nakatani T, Masuyama N, Taya S, Kumai M, Minaki Y, Hamaguchi A, Inoue YU, Inoue T, Miyashita S et al (2014) Temporal identity transition from Purkinje cell progenitors to GABAergic interneuron progenitors in the cerebellum. *Nat Commun* 5:3337
- Voogd J, Glickstein M (1998) The anatomy of the cerebellum. *Trends Neurosci* 21:370–375
- Wang VY, Rose MF, Zoghbi HY (2005) Math1 expression redefines the rhombic lip derivatives and reveals novel lineages within the brainstem and cerebellum. *Neuron* 48:31–43
- Wang SS, Kloth AD, Badura A (2014) The cerebellum, sensitive periods, and autism. *Neuron* 83:518–532
- Weisheit G, Gliem M, Endl E, Pfeffer PL, Busslinger M, Schilling K (2006) Postnatal development of the murine cerebellar cortex: formation and early dispersal of basket, stellate and Golgi neurons. *Eur J Neurosci* 24:466–478
- Wingate RJT (2001) The rhombic lip and early cerebellar development. *Curr Opin Neurobiol* 11:82–88
- Yamada M, Seto Y, Taya S, Owa T, Inoue YU, Inoue T, Kawaguchi Y, Nabeshima Y, Hoshino M (2014) Specification of spatial identities of cerebellar neuron progenitors by *ptf1a* and *atoh1* for proper production of GABAergic and glutamatergic neurons. *J Neurosci* 34:4786–4800
- Zordan P, Croci L, Hawkes R, Consalez GG (2008) Comparative analysis of proneural gene expression in the embryonic cerebellum. *Dev Dyn* 237:1726–1735